

SC35 Rabbit mAb
Catalog # AP76055**Specification**

SC35 Rabbit mAb - Product Information

Application	WB, IHC
Primary Accession	Q01130
Reactivity	Human, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	25476

SC35 Rabbit mAb - Additional Information**Gene ID** 6427**Other Names**
SRSF2**Dilution**
WB~~1/500-1/1000
IHC~~1/50-1/100**Format**
Liquid**SC35 Rabbit mAb - Protein Information****Name** SRSF2**Synonyms** SFRS2**Function**

Necessary for the splicing of pre-mRNA. It is required for formation of the earliest ATP-dependent splicing complex and interacts with spliceosomal components bound to both the 5'- and 3'-splice sites during spliceosome assembly. It also is required for ATP-dependent interactions of both U1 and U2 snRNPs with pre-mRNA. Interacts with other spliceosomal components, via the RS domains, to form a bridge between the 5'- and 3'-splice site binding components, U1 snRNP and U2AF. Binds to purine-rich RNA sequences, either 5'-AGSAGAGTA-3' (S=C or G) or 5'-GTTTCGAGTA-3'. Can bind to beta-globin mRNA and commit it to the splicing pathway. The phosphorylated form (by SRPK2) is required for cellular apoptosis in response to cisplatin treatment.

Cellular Location

Nucleus. Nucleus, nucleoplasm. Nucleus speckle. Note=Phosphorylation by SRPK2 provokes its redistribution from the nuclear speckle to nucleoplasm

SC35 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SC35 Rabbit mAb - Images

