

SF2 Rabbit mAb
Catalog # AP76073**Specification**

SF2 Rabbit mAb - Product Information

Application	WB, IHC, IF
Primary Accession	Q07955
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	27745

SF2 Rabbit mAb - Additional Information**Gene ID** 6426**Other Names**
SRSF1**Dilution**
WB~~1/500-1/1000
IHC~~1/50-1/100
IF~~1/50-1/200**Format**
Liquid**SF2 Rabbit mAb - Protein Information****Name** SRSF1 ([HGNC:10780](#))**Synonyms** ASF, SF2, SF2P33, SFRS1**Function**

Plays a role in preventing exon skipping, ensuring the accuracy of splicing and regulating alternative splicing. Interacts with other spliceosomal components, via the RS domains, to form a bridge between the 5'- and 3'-splice site binding components, U1 snRNP and U2AF. Can stimulate binding of U1 snRNP to a 5'-splice site- containing pre-mRNA. Binds to purine-rich RNA sequences, either the octamer, 5'-RGAAGAAC-3' (r=A or G) or the decamers, AGGACAGAGC/AGGACGAAGC. Binds preferentially to the 5'-CGAGGCG-3' motif in vitro. Three copies of the octamer constitute a powerful splicing enhancer in vitro, the ASF/SF2 splicing enhancer (ASE) which can specifically activate ASE-dependent splicing. Isoform ASF-2 and isoform ASF-3 act as splicing repressors. May function as export adapter involved in mRNA nuclear export through the TAP/NXF1 pathway.

Cellular Location

Cytoplasm. Nucleus speckle. Note=In nuclear speckles. Shuttles between the nucleus and the cytoplasm (PubMed:12215544, PubMed:20308322, PubMed:24449914, PubMed:9420331). Nuclear import is mediated via interaction with TNPO3 (PubMed:24449914).

SF2 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SF2 Rabbit mAb - Images



