

**TDP 43 Rabbit mAb**  
Catalog # AP76163**Specification****TDP 43 Rabbit mAb - Product Information**

Application	WB, IHC, IF
Primary Accession	<a href="#">Q13148</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	44740

**TDP 43 Rabbit mAb - Additional Information**

Gene ID 23435

**Other Names**

TARDBP

**Dilution**

WB~~1/500-1/1000

IHC~~1/50-1/100

IF~~1/50-1/200

**Format**

Liquid

**TDP 43 Rabbit mAb - Protein Information****Name** TARDBP {ECO:0000303|PubMed:18396105, ECO:0000312|HGNC:HGNC:11571}**Function**

RNA-binding protein that is involved in various steps of RNA biogenesis and processing (PubMed:<a href="http://www.uniprot.org/citations/23519609" target="\_blank">23519609</a>). Preferentially binds, via its two RNA recognition motifs RRM1 and RRM2, to GU-repeats on RNA molecules predominantly localized within long introns and in the 3'UTR of mRNAs (PubMed:<a href="http://www.uniprot.org/citations/23519609" target="\_blank">23519609</a>, PubMed:<a href="http://www.uniprot.org/citations/24240615" target="\_blank">24240615</a>, PubMed:<a href="http://www.uniprot.org/citations/24464995" target="\_blank">24464995</a>). In turn, regulates the splicing of many non-coding and protein-coding RNAs including proteins involved in neuronal survival, as well as mRNAs that encode proteins relevant for neurodegenerative diseases (PubMed:<a href="http://www.uniprot.org/citations/21358640" target="\_blank">21358640</a>, PubMed:<a href="http://www.uniprot.org/citations/29438978" target="\_blank">29438978</a>). Plays a role in maintaining mitochondrial homeostasis by regulating the processing of mitochondrial transcripts (PubMed:<a href="http://www.uniprot.org/citations/28794432" target="\_blank">28794432</a>). Regulates also mRNA stability by recruiting CNOT7/CAF1 deadenylase on mRNA 3'UTR leading to poly(A) tail deadenylation and thus shortening (PubMed:<a href="http://www.uniprot.org/citations/30520513" target="\_blank">30520513</a>).

In response to oxidative insult, associates with stalled ribosomes localized to stress granules (SGs) and contributes to cell survival (PubMed:[19765185](http://www.uniprot.org/citations/19765185), PubMed:[23398327](http://www.uniprot.org/citations/23398327)). Participates also in the normal skeletal muscle formation and regeneration, forming cytoplasmic myo-granules and binding mRNAs that encode sarcomeric proteins (PubMed:[30464263](http://www.uniprot.org/citations/30464263)). Plays a role in the maintenance of the circadian clock periodicity via stabilization of the CRY1 and CRY2 proteins in a FBXL3-dependent manner (PubMed:[27123980](http://www.uniprot.org/citations/27123980)). Negatively regulates the expression of CDK6 (PubMed:[19760257](http://www.uniprot.org/citations/19760257)). Regulates the expression of HDAC6, ATG7 and VCP in a PPIA/CYPA-dependent manner (PubMed:[25678563](http://www.uniprot.org/citations/25678563)).

#### Cellular Location

Nucleus. Cytoplasm. Cytoplasm, Stress granule Mitochondrion. Note=Continuously travels in and out of the nucleus (PubMed:18957508). Localizes to stress granules in response to oxidative stress (PubMed:19765185). A small subset localizes in mitochondria (PubMed:28794432).

#### Tissue Location

Ubiquitously expressed. In particular, expression is high in pancreas, placenta, lung, genital tract and spleen

#### TDP 43 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### TDP 43 Rabbit mAb - Images



