

**TRAF6 Rabbit mAb**  
Catalog # AP76191**Specification**

---

**TRAF6 Rabbit mAb - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">Q9Y4K3</a>
Reactivity	<b>Human, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Monoclonal Antibody</b>
Calculated MW	<b>59573</b>

**TRAF6 Rabbit mAb - Additional Information****Gene ID** 7189**Other Names**

TRAF6

**Dilution**

WB~~1/500-1/1000

**Format**

Liquid

**TRAF6 Rabbit mAb - Protein Information****Name** TRAF6**Synonyms** RNF85**Function**

E3 ubiquitin ligase that, together with UBE2N and UBE2V1, mediates the synthesis of 'Lys-63'-linked-polyubiquitin chains conjugated to proteins, such as ECSIT, IKBKG, IRAK1, AKT1 and AKT2 (PubMed: <a href="http://www.uniprot.org/citations/11057907" target="\_blank">11057907</a>, PubMed: <a href="http://www.uniprot.org/citations/18347055" target="\_blank">18347055</a>, PubMed: <a href="http://www.uniprot.org/citations/19465916" target="\_blank">19465916</a>, PubMed: <a href="http://www.uniprot.org/citations/19713527" target="\_blank">19713527</a>, PubMed: <a href="http://www.uniprot.org/citations/31620128" target="\_blank">31620128</a>). Also mediates ubiquitination of free/unanchored polyubiquitin chain that leads to MAP3K7 activation (PubMed: <a href="http://www.uniprot.org/citations/19675569" target="\_blank">19675569</a>). Leads to the activation of NF-kappa-B and JUN (PubMed: <a href="http://www.uniprot.org/citations/16378096" target="\_blank">16378096</a>, PubMed: <a href="http://www.uniprot.org/citations/17135271" target="\_blank">17135271</a>, PubMed: <a href="http://www.uniprot.org/citations/17703191" target="\_blank">17703191</a>). Seems to also play a role in dendritic cells (DCs) maturation and/or activation (By similarity). Represses c-Myb-mediated transactivation, in B-lymphocytes (PubMed: <a href="http://www.uniprot.org/citations/18093978" target="\_blank">18093978</a>),

PubMed:<a href="http://www.uniprot.org/citations/18758450" target="\_blank">18758450</a>). Adapter protein that seems to play a role in signal transduction initiated via TNF receptor, IL-1 receptor and IL-17 receptor (PubMed:<a href="http://www.uniprot.org/citations/12140561" target="\_blank">12140561</a>, PubMed:<a href="http://www.uniprot.org/citations/19825828" target="\_blank">19825828</a>, PubMed:<a href="http://www.uniprot.org/citations/8837778" target="\_blank">8837778</a>). Regulates osteoclast differentiation by mediating the activation of adapter protein complex 1 (AP-1) and NF-kappa-B, in response to RANK-L stimulation (By similarity). Together with MAP3K8, mediates CD40 signals that activate ERK in B-cells and macrophages, and thus may play a role in the regulation of immunoglobulin production (By similarity). Participates also in the TCR signaling by ubiquitinating LAT (PubMed:<a href="http://www.uniprot.org/citations/23514740" target="\_blank">23514740</a>, PubMed:<a href="http://www.uniprot.org/citations/25907557" target="\_blank">25907557</a>).

### Cellular Location

Cytoplasm. Cytoplasm, cell cortex. Nucleus. Lipid droplet {ECO:0000250|UniProtKB:P70196}. Note=Found in the nuclei of some aggressive B-cell lymphoma cell lines as well as in the nuclei of both resting and activated T- and B-lymphocytes. Found in punctate nuclear body protein complexes. Ubiquitination may occur in the cytoplasm and sumoylation in the nucleus. RSAD2/viperin recruits it to the lipid droplet (By similarity).

### Tissue Location

Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas

### TRAF6 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### TRAF6 Rabbit mAb - Images



