

TRAF6 Rabbit mAb
Catalog # AP76191**Specification****TRAF6 Rabbit mAb - Product Information**

Application	WB
Primary Accession	Q9Y4K3
Reactivity	Human, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	59573

TRAF6 Rabbit mAb - Additional Information

Gene ID 7189

Other Names

TRAF6

Dilution

WB~~1/500-1/1000

Format

Liquid

TRAF6 Rabbit mAb - Protein Information

Name TRAF6

Synonyms RNF85

Function

E3 ubiquitin ligase that, together with UBE2N and UBE2V1, mediates the synthesis of 'Lys-63'-linked-polyubiquitin chains conjugated to proteins, such as ECSIT, IKBKG, IRAK1, AKT1 and AKT2 (PubMed: [11057907](http://www.uniprot.org/citations/11057907) target="_blank">11057907, PubMed: [18347055](http://www.uniprot.org/citations/18347055) target="_blank">18347055, PubMed: [19465916](http://www.uniprot.org/citations/19465916) target="_blank">19465916, PubMed: [19713527](http://www.uniprot.org/citations/19713527) target="_blank">19713527, PubMed: [27746020](http://www.uniprot.org/citations/27746020) target="_blank">27746020, PubMed: [31620128](http://www.uniprot.org/citations/31620128) target="_blank">31620128). Also mediates ubiquitination of free/unanchored polyubiquitin chain that leads to MAP3K7 activation (PubMed: [19675569](http://www.uniprot.org/citations/19675569) target="_blank">19675569). Leads to the activation of NF-kappa-B and JUN (PubMed: [16378096](http://www.uniprot.org/citations/16378096) target="_blank">16378096, PubMed: [17135271](http://www.uniprot.org/citations/17135271) target="_blank">17135271, PubMed: [17703191](http://www.uniprot.org/citations/17703191) target="_blank">17703191). Seems to also play a role in dendritic cells (DCs) maturation and/or activation (By similarity). Represses c-Myb-mediated transactivation, in B-lymphocytes

(PubMed:18093978, PubMed:18758450). Adapter protein that seems to play a role in signal transduction initiated via TNF receptor, IL-1 receptor and IL-17 receptor (PubMed:12140561, PubMed:19825828, PubMed:8837778). Regulates osteoclast differentiation by mediating the activation of adapter protein complex 1 (AP-1) and NF-kappa-B, in response to RANK-L stimulation (By similarity). Together with MAP3K8, mediates CD40 signals that activate ERK in B-cells and macrophages, and thus may play a role in the regulation of immunoglobulin production (By similarity). Acts as a regulator of the JNK and NF-kappa-B signaling pathways by initiating assembly of heterotypic 'Lys-63'-/'Lys-48'-linked branched ubiquitin chains that are then recognized by TAB2: TRAF6 catalyzes initial 'Lys-63'-linked-polyubiquitin chains that are then branched via 'Lys-48'-linked polyubiquitin by HUWE1 (PubMed:27746020). 'Lys-63'-/'Lys-48'-linked branched ubiquitin chains protect 'Lys-63'- linkages from CYLD deubiquitination (PubMed:27746020). Participates also in the TCR signaling by ubiquitinating LAT (PubMed:23514740, PubMed:25907557).

Cellular Location

Cytoplasm. Cytoplasm, cell cortex. Nucleus. Lipid droplet {ECO:0000250|UniProtKB:P70196}. Note=Found in the nuclei of some aggressive B-cell lymphoma cell lines as well as in the nuclei of both resting and activated T- and B-lymphocytes. Found in punctate nuclear body protein complexes. Ubiquitination may occur in the cytoplasm and sumoylation in the nucleus. RSAD2/viperin recruits it to the lipid droplet (By similarity).

Tissue Location

Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas

TRAF6 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TRAF6 Rabbit mAb - Images



