

TRAF6 Rabbit mAb
Catalog # AP76192**Specification**

TRAF6 Rabbit mAb - Product Information

Application	WB
Primary Accession	Q9Y4K3
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	59573

TRAF6 Rabbit mAb - Additional Information**Gene ID** 7189**Other Names**
TRAF6**Dilution**
WB~~1/500-1/1000**Format**
Liquid**TRAF6 Rabbit mAb - Protein Information****Name** TRAF6**Synonyms** RNF85**Function**

E3 ubiquitin ligase that, together with UBE2N and UBE2V1, mediates the synthesis of 'Lys-63'-linked-polyubiquitin chains conjugated to proteins, such as ECSIT, IKBKG, IRAK1, AKT1 and AKT2 (PubMed: [11057907](http://www.uniprot.org/citations/11057907) target="_blank">11057907, PubMed: [18347055](http://www.uniprot.org/citations/18347055) target="_blank">18347055, PubMed: [19465916](http://www.uniprot.org/citations/19465916) target="_blank">19465916, PubMed: [19713527](http://www.uniprot.org/citations/19713527) target="_blank">19713527, PubMed: [31620128](http://www.uniprot.org/citations/31620128) target="_blank">31620128). Also mediates ubiquitination of free/unanchored polyubiquitin chain that leads to MAP3K7 activation (PubMed: [19675569](http://www.uniprot.org/citations/19675569) target="_blank">19675569). Leads to the activation of NF-kappa-B and JUN (PubMed: [16378096](http://www.uniprot.org/citations/16378096) target="_blank">16378096, PubMed: [17135271](http://www.uniprot.org/citations/17135271) target="_blank">17135271, PubMed: [17703191](http://www.uniprot.org/citations/17703191) target="_blank">17703191). Seems to also play a role in dendritic cells (DCs) maturation and/or activation (By similarity). Represses c-Myb-mediated transactivation, in B-lymphocytes (PubMed: [18093978](http://www.uniprot.org/citations/18093978) target="_blank">18093978, PubMed: [18758450](http://www.uniprot.org/citations/18758450) target="_blank">18758450).

Adapter protein that seems to play a role in signal transduction initiated via TNF receptor, IL-1 receptor and IL-17 receptor (PubMed:12140561, PubMed:19825828, PubMed:8837778). Regulates osteoclast differentiation by mediating the activation of adapter protein complex 1 (AP-1) and NF-kappa-B, in response to RANK-L stimulation (By similarity). Together with MAP3K8, mediates CD40 signals that activate ERK in B-cells and macrophages, and thus may play a role in the regulation of immunoglobulin production (By similarity). Participates also in the TCR signaling by ubiquitinating LAT (PubMed:23514740, PubMed:25907557).

Cellular Location

Cytoplasm. Cytoplasm, cell cortex. Nucleus. Lipid droplet {ECO:0000250|UniProtKB:P70196}. Note=Found in the nuclei of some aggressive B-cell lymphoma cell lines as well as in the nuclei of both resting and activated T- and B-lymphocytes. Found in punctate nuclear body protein complexes. Ubiquitination may occur in the cytoplasm and sumoylation in the nucleus. RSAD2/viperin recruits it to the lipid droplet (By similarity).

Tissue Location

Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas

TRAF6 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TRAF6 Rabbit mAb - Images



