

ATF6 Rabbit mAb
Catalog # AP76397**Specification**

ATF6 Rabbit mAb - Product Information

Application	WB, IHC
Primary Accession	P18850
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	74585

ATF6 Rabbit mAb - Additional Information**Gene ID** 22926**Other Names**
ATF6**Dilution**
WB~~1/500-1/1000
IHC~~1/50-1/100**Format**
Liquid**ATF6 Rabbit mAb - Protein Information****Name** ATF6**Function**

[Cyclic AMP-dependent transcription factor ATF-6 alpha]: Precursor of the transcription factor form (Processed cyclic AMP- dependent transcription factor ATF-6 alpha), which is embedded in the endoplasmic reticulum membrane (PubMed: [10564271](http://www.uniprot.org/citations/10564271), PubMed: [11158310](http://www.uniprot.org/citations/11158310), PubMed: [11779464](http://www.uniprot.org/citations/11779464)). Endoplasmic reticulum stress promotes processing of this form, releasing the transcription factor form that translocates into the nucleus, where it activates transcription of genes involved in the unfolded protein response (UPR) (PubMed: [10564271](http://www.uniprot.org/citations/10564271), PubMed: [11158310](http://www.uniprot.org/citations/11158310), PubMed: [11779464](http://www.uniprot.org/citations/11779464)).

Cellular Location

Endoplasmic reticulum membrane; Single-pass type II membrane protein. Golgi apparatus membrane; Single-pass type II membrane protein. Note=Translocates from the endoplasmic reticulum to the Golgi, where it is processed.

Tissue Location
Ubiquitous..

ATF6 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ATF6 Rabbit mAb - Images

