

BMI1 Rabbit mAb
Catalog # AP76410**Specification**

BMI1 Rabbit mAb - Product Information

Application	WB, IF, IP
Primary Accession	P35226
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	36949

BMI1 Rabbit mAb - Additional Information**Gene ID** 100532731;648**Other Names**

BMI1

Dilution

WB~~1/500-1/1000

IF~~1/50-1/200

IP~~1/20

Format

Liquid

BMI1 Rabbit mAb - Protein Information**Name** BMI1**Synonyms** PCGF4, RNF51**Function**

Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility (PubMed: 15386022, PubMed: 16359901, PubMed: 16714294, PubMed: 21772249, PubMed: 25355358, PubMed: 26151332, PubMed: 27827373). The complex composed of RNF2, UB2D3 and BMI1 binds nucleosomes, and has activity only with nucleosomal histone H2A (PubMed: 21772249, PubMed: 21772249, PubMed: 21772249).

[25355358](http://www.uniprot.org/citations/25355358)). In the PRC1-like complex, regulates the E3 ubiquitin-protein ligase activity of RNF2/RING2 (PubMed:[15386022](http://www.uniprot.org/citations/15386022), PubMed:[21772249](http://www.uniprot.org/citations/21772249), PubMed:[26151332](http://www.uniprot.org/citations/26151332)).

Cellular Location

Nucleus. Cytoplasm

BMI1 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BMI1 Rabbit mAb - Images



