

**Keap1 Rabbit mAb**  
Catalog # AP76559**Specification****Keap1 Rabbit mAb - Product Information**

Application	<b>WB, IHC</b>
Primary Accession	<a href="#">Q14145</a>
Reactivity	<b>Human, Mouse</b>
Host	<b>Rabbit</b>
Clonality	<b>Monoclonal Antibody</b>
Calculated MW	<b>69666</b>

**Keap1 Rabbit mAb - Additional Information****Gene ID** 9817**Other Names**

KEAP1

**Dilution**

WB~~1/500-1/1000

IHC~~1/50-1/100

**Format**

Liquid

**Keap1 Rabbit mAb - Protein Information****Name** KEAP1 {ECO:0000303|PubMed:14585973, ECO:0000312|HGNC:HGNC:23177}**Function**

Substrate-specific adapter of a BCR (BTB-CUL3-RBX1) E3 ubiquitin ligase complex that regulates the response to oxidative stress by targeting NFE2L2/NRF2 for ubiquitination (PubMed: [14585973](http://www.uniprot.org/citations/14585973), PubMed: [15379550](http://www.uniprot.org/citations/15379550), PubMed: [15572695](http://www.uniprot.org/citations/15572695), PubMed: [15601839](http://www.uniprot.org/citations/15601839), PubMed: [15983046](http://www.uniprot.org/citations/15983046), PubMed: [37339955](http://www.uniprot.org/citations/37339955)). KEAP1 acts as a key sensor of oxidative and electrophilic stress: in normal conditions, the BCR(KEAP1) complex mediates ubiquitination and degradation of NFE2L2/NRF2, a transcription factor regulating expression of many cytoprotective genes (PubMed: [15601839](http://www.uniprot.org/citations/15601839), PubMed: [16006525](http://www.uniprot.org/citations/16006525)). In response to oxidative stress, different electrophile metabolites trigger non-enzymatic covalent modifications of highly reactive cysteine residues in KEAP1, leading to inactivate the ubiquitin ligase activity of the BCR(KEAP1) complex, promoting NFE2L2/NRF2 nuclear accumulation and expression of phase II detoxifying enzymes (PubMed: [16006525](http://www.uniprot.org/citations/16006525))

target="\_blank">16006525</a>, PubMed:<a href="http://www.uniprot.org/citations/17127771" target="\_blank">17127771</a>, PubMed:<a href="http://www.uniprot.org/citations/18251510" target="\_blank">18251510</a>, PubMed:<a href="http://www.uniprot.org/citations/19489739" target="\_blank">19489739</a>, PubMed:<a href="http://www.uniprot.org/citations/29590092" target="\_blank">29590092</a>). In response to selective autophagy, KEAP1 is sequestered in inclusion bodies following its interaction with SQSTM1/p62, leading to inactivation of the BCR(KEAP1) complex and activation of NFE2L2/NRF2 (PubMed:<a href="http://www.uniprot.org/citations/20452972" target="\_blank">20452972</a>). The BCR(KEAP1) complex also mediates ubiquitination of SQSTM1/p62, increasing SQSTM1/p62 sequestering activity and degradation (PubMed:<a href="http://www.uniprot.org/citations/28380357" target="\_blank">28380357</a>). The BCR(KEAP1) complex also targets BPTF and PGAM5 for ubiquitination and degradation by the proteasome (PubMed:<a href="http://www.uniprot.org/citations/15379550" target="\_blank">15379550</a>, PubMed:<a href="http://www.uniprot.org/citations/17046835" target="\_blank">17046835</a>).

#### **Cellular Location**

Cytoplasm. Nucleus. Note=Mainly cytoplasmic (PubMed:15601839). In response to selective autophagy, relocates to inclusion bodies following interaction with SQSTM1/p62 (PubMed:20452972).

#### **Tissue Location**

Broadly expressed, with highest levels in skeletal muscle.

#### **Keap1 Rabbit mAb - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **Keap1 Rabbit mAb - Images**



