

Ku70 Rabbit mAb
Catalog # AP76564**Specification****Ku70 Rabbit mAb - Product Information**

Application	WB, IHC, IF
Primary Accession	P12956
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	69843

Ku70 Rabbit mAb - Additional Information

Gene ID 2547

Other Names

XRCC6

Dilution

WB~~1/500-1/1000

IHC~~1/50-1/100

IF~~1/50-1/200

Format

Liquid

Ku70 Rabbit mAb - Protein Information

Name XRCC6

Synonyms G22P1

Function

Single-stranded DNA-dependent ATP-dependent helicase that plays a key role in DNA non-homologous end joining (NHEJ) by recruiting DNA-PK to DNA (PubMed: [11493912](http://www.uniprot.org/citations/11493912), PubMed: [12145306](http://www.uniprot.org/citations/12145306), PubMed: [20493174](http://www.uniprot.org/citations/20493174), PubMed: [2466842](http://www.uniprot.org/citations/2466842), PubMed: [7957065](http://www.uniprot.org/citations/7957065), PubMed: [8621488](http://www.uniprot.org/citations/8621488), PubMed: [9742108](http://www.uniprot.org/citations/9742108)). Required for double-strand break repair and V(D)J recombination (PubMed: [11493912](http://www.uniprot.org/citations/11493912), PubMed: [12145306](http://www.uniprot.org/citations/12145306), PubMed: [20493174](http://www.uniprot.org/citations/20493174), PubMed: [2466842](http://www.uniprot.org/citations/2466842), PubMed: [7957065](http://www.uniprot.org/citations/7957065), PubMed: [8621488](http://www.uniprot.org/citations/8621488), PubMed: [9742108](http://www.uniprot.org/citations/9742108)).

href="http://www.uniprot.org/citations/7957065" target="_blank">7957065, PubMed:8621488, PubMed:9742108). Also has a role in chromosome translocation (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065, PubMed:8621488, PubMed:9742108). Has a role in chromosome translocation (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065, PubMed:8621488, PubMed:9742108). The DNA helicase II complex binds preferentially to fork-like ends of double-stranded DNA in a cell cycle-dependent manner (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065, PubMed:8621488, PubMed:9742108). It works in the 3'-5' direction (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065, PubMed:8621488, PubMed:9742108). During NHEJ, the XRCC5-XRCC6 dimer performs the recognition step: it recognizes and binds to the broken ends of the DNA and protects them from further resection (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065, PubMed:8621488, PubMed:9742108). Binding to DNA may be mediated by XRCC6 (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065, PubMed:8621488, PubMed:9742108). The XRCC5-XRCC6 dimer acts as a regulatory subunit of the DNA-dependent protein kinase complex DNA-PK by increasing the affinity of the catalytic subunit PRKDC to DNA by 100-fold (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065

target="_blank">7957065, PubMed:8621488, PubMed:9742108). The XRCC5-XRCC6 dimer is probably involved in stabilizing broken DNA ends and bringing them together (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065, PubMed:8621488, PubMed:9742108). The assembly of the DNA-PK complex to DNA ends is required for the NHEJ ligation step (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065, PubMed:8621488, PubMed:9742108). Probably also acts as a 5'-deoxyribose-5-phosphate lyase (5'-dRP lyase), by catalyzing the beta-elimination of the 5' deoxyribose-5-phosphate at an abasic site near double-strand breaks (PubMed:20383123). 5'-dRP lyase activity allows to 'clean' the termini of abasic sites, a class of nucleotide damage commonly associated with strand breaks, before such broken ends can be joined (PubMed:20383123). The XRCC5-XRCC6 dimer together with APEX1 acts as a negative regulator of transcription (PubMed:8621488). In association with NAA15, the XRCC5-XRCC6 dimer binds to the osteocalcin promoter and activates osteocalcin expression (PubMed:12145306). Plays a role in the regulation of DNA virus-mediated innate immune response by assembling into the HDP-RNP complex, a complex that serves as a platform for IRF3 phosphorylation and subsequent innate immune response activation through the cGAS-STING pathway (PubMed:28712728).

Cellular Location

Nucleus. Chromosome

Ku70 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Ku70 Rabbit mAb - Images



