

**PIM2 Rabbit mAb**  
Catalog # AP76657**Specification**

---

**PIM2 Rabbit mAb - Product Information**

|                   |                            |
|-------------------|----------------------------|
| Application       | <b>WB</b>                  |
| Primary Accession | <a href="#">Q9P1W9</a>     |
| Reactivity        | <b>Human</b>               |
| Host              | <b>Rabbit</b>              |
| Clonality         | <b>Monoclonal Antibody</b> |
| Calculated MW     | <b>34190</b>               |

**PIM2 Rabbit mAb - Additional Information****Gene ID** 11040**Other Names**  
PIM2**Dilution**  
WB~~1/500-1/1000**Format**  
Liquid**PIM2 Rabbit mAb - Protein Information****Name** PIM2**Function**

Proto-oncogene with serine/threonine kinase activity involved in cell survival and cell proliferation. Exerts its oncogenic activity through: the regulation of MYC transcriptional activity, the regulation of cell cycle progression, the regulation of cap-dependent protein translation and through survival signaling by phosphorylation of a pro- apoptotic protein, BAD. Phosphorylation of MYC leads to an increase of MYC protein stability and thereby an increase transcriptional activity. The stabilization of MYC exerted by PIM2 might explain partly the strong synergism between these 2 oncogenes in tumorigenesis. Regulates cap-dependent protein translation in a mammalian target of rapamycin complex 1 (mTORC1)-independent manner and in parallel to the PI3K-Akt pathway. Mediates survival signaling through phosphorylation of BAD, which induces release of the anti-apoptotic protein Bcl-X(L)/BCL2L1. Promotes cell survival in response to a variety of proliferative signals via positive regulation of the I-kappa-B kinase/NF-kappa-B cascade; this process requires phosphorylation of MAP3K8/COT. Promotes growth factor-independent proliferation by phosphorylation of cell cycle factors such as CDKN1A and CDKN1B. Involved in the positive regulation of chondrocyte survival and autophagy in the epiphyseal growth plate.

**Tissue Location**

Highly expressed in hematopoietic tissues, in leukemic and lymphoma cell lines, testis, small intestine, colon and colorectal adenocarcinoma cells. Weakly expressed in normal liver, but highly

expressed in hepatocellular carcinoma tissues

### PIM2 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### PIM2 Rabbit mAb - Images

