

Rab29 Rabbit mAb
Catalog # AP76681**Specification**

Rab29 Rabbit mAb - Product Information

Application	WB
Primary Accession	O14966
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	23155

Rab29 Rabbit mAb - Additional Information**Gene ID** 8934**Other Names**

RAB29

Dilution

WB~~1/500-1/1000

Format

Liquid

Rab29 Rabbit mAb - Protein Information**Name** RAB29**Synonyms** RAB7L1**Function**

The small GTPases Rab are key regulators in vesicle trafficking (PubMed: [24788816](http://www.uniprot.org/citations/24788816)). Essential for maintaining the integrity of the endosome-trans-Golgi network structure (By similarity). Together with LRRK2, plays a role in the retrograde trafficking pathway for recycling proteins, such as mannose 6 phosphate receptor (M6PR), between lysosomes and the Golgi apparatus in a retromer-dependent manner (PubMed: [24788816](http://www.uniprot.org/citations/24788816)). Recruits LRRK2 to the Golgi complex and stimulates LRRK2 kinase activity (PubMed: [29212815](http://www.uniprot.org/citations/29212815), PubMed: [38127736](http://www.uniprot.org/citations/38127736)). Stimulates phosphorylation of RAB10 'Thr-73' by LRRK2 (PubMed: [38127736](http://www.uniprot.org/citations/38127736)). Regulates neuronal process morphology in the intact central nervous system (CNS) (By similarity). May play a role in the formation of typhoid toxin transport intermediates during Salmonella enterica serovar Typhi (S.Typhi) epithelial cell infection (PubMed: [22042847](http://www.uniprot.org/citations/22042847)).

Cellular Location

Cell membrane; Lipid-anchor; Cytoplasmic side. Cytoplasm. Cytoplasm, perinuclear region. Golgi apparatus. Golgi apparatus membrane. Golgi apparatus, trans-Golgi network. Vacuole. Cytoplasm, cytoskeleton. Note=Colocalizes with LRRK2 along tubular structures emerging from Golgi apparatus (PubMed:29212815) Colocalizes with GM130 at the Golgi apparatus (PubMed:22042847) Colocalizes with dynamic tubules emerging from and retracting to the Golgi apparatus (PubMed:22042847, PubMed:38127736). Colocalizes with TGN46 at the trans-Golgi network (TGN) (PubMed:24788816). In *Salmonella enterica* serovar Typhi (S.Typhi) infected epithelial cells, is recruited and colocalized with both S.Typhi-containing vacuoles and dynamic tubules as well as those emerging from the vacuole toward the cell periphery (PubMed:22042847).

Tissue Location

Ubiquitous..

Rab29 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Rab29 Rabbit mAb - Images

