

Ubiquitin 60S Ribosomal Protein L40 Rabbit mAb
Catalog # AP76753**Specification**

Ubiquitin 60S Ribosomal Protein L40 Rabbit mAb - Product Information

Application	WB, IF
Primary Accession	P62987
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	14728

Ubiquitin 60S Ribosomal Protein L40 Rabbit mAb - Additional Information**Gene ID** 7311**Other Names**
UBA52**Dilution**
WB~~1/500-1/1000
IF~~1/50-1/200**Format**
Liquid**Ubiquitin 60S Ribosomal Protein L40 Rabbit mAb - Protein Information****Name** UBA52**Synonyms** UBCEP2**Function**

[Ubiquitin]: Exists either covalently attached to another protein, or free (unanchored). When covalently bound, it is conjugated to target proteins via an isopeptide bond either as a monomer (monoubiquitin), a polymer linked via different Lys residues of the ubiquitin (polyubiquitin chains) or a linear polymer linked via the initiator Met of the ubiquitin (linear polyubiquitin chains). Polyubiquitin chains, when attached to a target protein, have different functions depending on the Lys residue of the ubiquitin that is linked: Lys-6-linked may be involved in DNA repair; Lys-11-linked is involved in ERAD (endoplasmic reticulum-associated degradation) and in cell-cycle regulation; Lys-29-linked is involved in proteotoxic stress response and cell cycle; Lys-33-linked is involved in kinase modification; Lys-48-linked is involved in protein degradation via the proteasome; Lys-63-linked is involved in endocytosis, DNA-damage responses as well as in signaling processes leading to activation of the transcription factor NF-kappa-B. Linear polymer chains formed via attachment by the initiator Met lead to cell signaling. Ubiquitin is usually conjugated to Lys residues of target proteins, however, in rare cases, conjugation to Cys or Ser residues has been observed. When polyubiquitin is free (unanchored-polyubiquitin), it also has distinct roles, such as in activation of protein kinases, and in signaling.

Cellular Location

[Ubiquitin]: Cytoplasm. Nucleus

Ubiquitin 60S Ribosomal Protein L40 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Ubiquitin 60S Ribosomal Protein L40 Rabbit mAb - Images



