

**Ubiquitin Protein Ligase E3A Rabbit mAb**  
Catalog # AP76758**Specification****Ubiquitin Protein Ligase E3A Rabbit mAb - Product Information**

Application	WB, IF, IP
Primary Accession	<a href="#">Q05086</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	100688

**Ubiquitin Protein Ligase E3A Rabbit mAb - Additional Information**

Gene ID 7337

**Other Names**

UBE3A

**Dilution**

WB~~1/500-1/1000

IF~~1/50-1/200

IP~~1/20

**Format**

Liquid

**Ubiquitin Protein Ligase E3A Rabbit mAb - Protein Information**Name UBE3A ([HGNC:12496](#))**Function**

E3 ubiquitin-protein ligase which accepts ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and transfers it to its substrates (PubMed:[10373495](http://www.uniprot.org/citations/10373495), PubMed:[16772533](http://www.uniprot.org/citations/16772533), PubMed:[19204938](http://www.uniprot.org/citations/19204938), PubMed:[19233847](http://www.uniprot.org/citations/19233847), PubMed:[19325566](http://www.uniprot.org/citations/19325566), PubMed:[19591933](http://www.uniprot.org/citations/19591933), PubMed:[22645313](http://www.uniprot.org/citations/22645313), PubMed:[24273172](http://www.uniprot.org/citations/24273172), PubMed:[24728990](http://www.uniprot.org/citations/24728990), PubMed:[30020076](http://www.uniprot.org/citations/30020076)). Several substrates have been identified including the BMAL1, ARC, LAMTOR1, RAD23A and RAD23B, MCM7 (which is involved in DNA replication), annexin A1, the PML tumor suppressor, and the cell cycle regulator CDKN1B (PubMed:[10373495](http://www.uniprot.org/citations/10373495), PubMed:[19204938](http://www.uniprot.org/citations/19204938)

target="\_blank">19204938</a>, PubMed:<a href="http://www.uniprot.org/citations/19325566" target="\_blank">19325566</a>, PubMed:<a href="http://www.uniprot.org/citations/19591933" target="\_blank">19591933</a>, PubMed:<a href="http://www.uniprot.org/citations/22645313" target="\_blank">22645313</a>, PubMed:<a href="http://www.uniprot.org/citations/24728990" target="\_blank">24728990</a>, PubMed:<a href="http://www.uniprot.org/citations/30020076" target="\_blank">30020076</a>). Additionally, may function as a cellular quality control ubiquitin ligase by helping the degradation of the cytoplasmic misfolded proteins (PubMed:<a href="http://www.uniprot.org/citations/19233847" target="\_blank">19233847</a>). Finally, UBE3A also promotes its own degradation in vivo. Plays an important role in the regulation of the circadian clock: involved in the ubiquitination of the core clock component BMAL1, leading to its proteasomal degradation (PubMed:<a href="http://www.uniprot.org/citations/24728990" target="\_blank">24728990</a>). Acts as transcriptional coactivator of progesterone receptor PGR upon progesterone hormone activation (PubMed:<a href="http://www.uniprot.org/citations/16772533" target="\_blank">16772533</a>). Acts as a regulator of synaptic development by mediating ubiquitination and degradation of ARC (By similarity). Required for synaptic remodeling in neurons by mediating ubiquitination and degradation of LAMTOR1, thereby limiting mTORC1 signaling and activity-dependent synaptic remodeling (By similarity). Synergizes with WBP2 in enhancing PGR activity (PubMed:<a href="http://www.uniprot.org/citations/16772533" target="\_blank">16772533</a>).

#### Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:O08759}. Nucleus {ECO:0000250|UniProtKB:O08759}

#### Ubiquitin Protein Ligase E3A Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Ubiquitin Protein Ligase E3A Rabbit mAb - Images



