

Ki67 Rabbit mAb
Catalog # AP76932**Specification**

Ki67 Rabbit mAb - Product Information

Application	WB, IHC-P, FC, ICC
Primary Accession	P46013
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	358694

Ki67 Rabbit mAb - Additional Information

Gene ID 4288

Other Names

MKI67

Format

Liquid

Ki67 Rabbit mAb - Protein InformationName MKI67 ([HGNC:7107](#))**Function**

Protein that associates with the surface of mitotic chromosomes and acts both as a chromosome repellent during early mitosis and chromosome attractant during late mitosis (PubMed: [27362226](http://www.uniprot.org/citations/27362226), PubMed: [32879492](http://www.uniprot.org/citations/32879492), PubMed: [35513709](http://www.uniprot.org/citations/35513709), PubMed: [39153474](http://www.uniprot.org/citations/39153474)). Required to maintain individual mitotic chromosomes dispersed in the cytoplasm following nuclear envelope disassembly (PubMed: [27362226](http://www.uniprot.org/citations/27362226)). During early mitosis, relocalizes from nucleoli to the chromosome surface where it forms extended brush structures that cover a substantial fraction of the chromosome surface (PubMed: [27362226](http://www.uniprot.org/citations/27362226)). The MKI67 brush structure prevents chromosomes from collapsing into a single chromatin mass by forming a steric and electrostatic charge barrier: the protein has a high net electrical charge and acts as a surfactant, dispersing chromosomes and enabling independent chromosome motility (PubMed: [27362226](http://www.uniprot.org/citations/27362226)). During mitotic anaphase, the MKI67 brush structure collapses and MKI67 switches from a chromosome repellent to a chromosome attractant to promote chromosome clustering and facilitate the exclusion of large cytoplasmic particles from the future nuclear space (PubMed: [32879492](http://www.uniprot.org/citations/32879492), PubMed: [39153474](http://www.uniprot.org/citations/39153474)).

Mechanistically, dephosphorylation during mitotic exit and simultaneous exposure of a conserved basic patch induce the RNA-dependent formation of a liquid-like condensed phase on the chromosome surface, promoting coalescence of neighboring chromosome surfaces and clustering of chromosomes (PubMed:[39153474](http://www.uniprot.org/citations/39153474) target="_blank">39153474). Binds premature ribosomal RNAs during anaphase; promoting liquid-liquid phase separation (PubMed:[28935370](http://www.uniprot.org/citations/28935370) target="_blank">28935370), PubMed:[39153474](http://www.uniprot.org/citations/39153474) target="_blank">39153474). Binds DNA, with a preference for supercoiled DNA and AT-rich DNA (PubMed:[10878551](http://www.uniprot.org/citations/10878551) target="_blank">10878551). Does not contribute to the internal structure of mitotic chromosomes (By similarity). May play a role in chromatin organization; it is however unclear whether it plays a direct role in chromatin organization or whether it is an indirect consequence of its function in mitotic chromosome (PubMed:[24867636](http://www.uniprot.org/citations/24867636) target="_blank">24867636).

Cellular Location

Chromosome. Nucleus. Nucleus, nucleolus. Note=During early mitosis, relocates from nucleoli to the surface of the mitotic chromosome, the perichromosomal layer, and covers a substantial fraction of the mitotic chromosome surface (PubMed:27362226) Associates with satellite DNA in G1 phase (PubMed:9510506). Binds tightly to chromatin in interphase, chromatin-binding decreases in mitosis when it associates with the surface of the condensed chromosomes (PubMed:15896774, PubMed:22002106). Predominantly localized in the G1 phase in the perinucleolar region, in the later phases it is also detected throughout the nuclear interior, being predominantly localized in the nuclear matrix (PubMed:22002106)

Ki67 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Ki67 Rabbit mAb - Images