

VAMP8 Rabbit mAb

Catalog # AP76954

Specification

VAMP8 Rabbit mAb - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW

WB
O9BV40
Human, Mouse, Rat
Rabbit
Monoclonal Antibody
11438

VAMP8 Rabbit mAb - Additional Information

Gene ID 8673

Other Names VAMP8

Dilution WB~~1/500-1/1000

Format Liquid

VAMP8 Rabbit mAb - Protein Information

Name VAMP8 {ECO:0000303|PubMed:12130530}

Function

SNAREs, soluble N-ethylmaleimide-sensitive factor-attachment protein receptors, are essential proteins for fusion of cellular membranes. SNAREs localized on opposing membranes assemble to form a trans-SNARE complex, an extended, parallel four alpha-helical bundle that drives membrane fusion. VAMP8 is a SNARE involved in autophagy through the direct control of autophagosome membrane fusion with the lysososome membrane via its interaction with the STX17-SNAP29 binary t- SNARE complex (PubMed:<a

 $href="http://www.uniprot.org/citations/23217709" target="_blank">23217709, PubMed:25686604). Also required for dense-granule secretion in platelets (PubMed:<a$

href="http://www.uniprot.org/citations/12130530" target="_blank">12130530). Also plays a role in regulated enzyme secretion in pancreatic acinar cells (By similarity). Involved in the abscission of the midbody during cell division, which leads to completely separate daughter cells (By similarity). Involved in the homotypic fusion of early and late endosomes (By similarity). Participates also in the activation of type I interferon antiviral response through a TRIM6-dependent mechanism (PubMed:31694946).

Cellular Location





Lysosome membrane; Single-pass type IV membrane protein. Early endosome membrane; Single-pass type IV membrane protein. Late endosome membrane; Single-pass type IV membrane protein. Cell membrane {ECO:0000250|UniProtKB:070404}; Single-pass type IV membrane protein. Zymogen granule membrane {ECO:0000250|UniProtKB:070404}; Single-pass type IV membrane protein. Note=Perinuclear vesicular structures of the early and late endosomes, coated pits, and trans-Golgi (By similarity) Sub-tight junctional domain in retinal pigment epithelium cells Midbody region during cytokinesis. Lumenal oriented, apical membranes of nephric tubular cell (By similarity). Cycles through the apical but not through the basolateral plasma membrane (By similarity). Apical region of acinar cells; in zymogen granule membranes (By similarity) {ECO:0000250|UniProtKB:Q9WUF4}

Tissue Location Platelets..

VAMP8 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

VAMP8 Rabbit mAb - Images

