

# TRPM8 Rabbit mAb

**Catalog # AP76975** 

# **Specification**

#### TRPM8 Rabbit mAb - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW

WB
O7Z2W7
Human, Mouse
Rabbit
Monoclonal Antibody
127685

#### TRPM8 Rabbit mAb - Additional Information

Gene ID 79054

Other Names TRPM8

**Dilution** WB~~1/500-1/1000

Format Liquid

### **TRPM8 Rabbit mAb - Protein Information**

Name TRPM8

Synonyms LTRPC6, TRPP8

#### **Function**

Non-selective ion channel permeable to monovalent and divalent cations, including Na(+), K(+), and Ca(2+), with higher permeability for Ca(2+). Activated by multiple factors, such as temperature, voltage, pressure, and changes in osmolality. Activated by cool temperatures (<23-28 degrees Celsius) and by chemical ligands evoking a sensation of coolness, such as menthol and icilin therefore plays a central role in the detection of environmental cold temperatures (PubMed:<a href="http://www.uniprot.org/citations/15306801" target=" blank">15306801</a>, PubMed:<a href="http://www.uniprot.org/citations/15852009" target="blank">15852009</a>, PubMed:<a href="http://www.uniprot.org/citations/16174775" target="blank">16174775</a>, PubMed:<a href="http://www.uniprot.org/citations/25559186" target="blank">25559186</a>, PubMed:<a href="http://www.uniprot.org/citations/37857704" target="blank">37857704</a>). TRPM8 is a voltage-dependent channel; its activation by cold or chemical ligands shifts its voltage thresholds towards physiological membrane potentials, leading to the opening of the channel (PubMed:<a href="http://www.uniprot.org/citations/15306801" target=" blank">15306801</a>). In addition to its critical role in temperature sensing, regulates basal tear secretion by sensing evaporation-induced cooling and changes in osmolality (By similarity). May plays a role in prostate cancer cell migration (PubMed: <a



href="http://www.uniprot.org/citations/16174775" target="\_blank">16174775</a>, PubMed:<a href="http://www.uniprot.org/citations/25559186" target="\_blank">25559186</a>).

#### **Cellular Location**

Cell membrane; Multi-pass membrane protein. Membrane raft {ECO:0000250|UniProtKB:Q8R4D5}. Endoplasmic reticulum membrane. Note=Lipid raft association modulates TRPM8 channel activity (By similarity) Located in the endoplasmic reticulum in prostate cancer cells (PubMed:11325849, PubMed:16174775). {ECO:0000250|UniProtKB:Q8R4D5, ECO:0000269|PubMed:11325849, ECO:0000269|PubMed:16174775}

### **Tissue Location**

Expressed in prostate. Also expressed in prostate tumors and in non-prostatic primary tumors such as colon, lung, breast and skin tumors.

## TRPM8 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# TRPM8 Rabbit mAb - Images

