

MOV10 Rabbit mAb
Catalog # AP76996**Specification****MOV10 Rabbit mAb - Product Information**

Application	WB
Primary Accession	Q9HCE1
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	113671

MOV10 Rabbit mAb - Additional Information**Gene ID** 4343**Other Names**

MOV10

Dilution

WB~~1/500-1/1000

Format

Liquid

MOV10 Rabbit mAb - Protein Information**Name** MOV10 ([HGNC:7200](#))**Synonyms** KIAA1631**Function**

5' to 3' RNA helicase that is involved in a number of cellular roles ranging from mRNA metabolism and translation, modulation of viral infectivity, inhibition of retrotransposition, or regulation of synaptic transmission (PubMed:[23093941](http://www.uniprot.org/citations/23093941)). Plays an important role in innate antiviral immunity by promoting type I interferon production (PubMed:[27016603](http://www.uniprot.org/citations/27016603), PubMed:[27974568](http://www.uniprot.org/citations/27974568), PubMed:[35157734](http://www.uniprot.org/citations/35157734)). Mechanistically, specifically uses IKKepsilon/IKBKE as the mediator kinase for IRF3 activation (PubMed:[27016603](http://www.uniprot.org/citations/27016603), PubMed:[35157734](http://www.uniprot.org/citations/35157734)). Blocks HIV-1 virus replication at a post-entry step (PubMed:[20215113](http://www.uniprot.org/citations/20215113)). Counteracts HIV-1 Vif-mediated degradation of APOBEC3G through its helicase activity by interfering with the ubiquitin-proteasome pathway (PubMed:[29258557](http://www.uniprot.org/citations/29258557)). Inhibits also hepatitis B virus/HBV replication by interacting with

HBV RNA and thereby inhibiting the early step of viral reverse transcription (PubMed:31722967). Contributes to UPF1 mRNA target degradation by translocation along 3' UTRs (PubMed:24726324). Required for microRNA (miRNA)-mediated gene silencing by the RNA-induced silencing complex (RISC). Required for both miRNA-mediated translational repression and miRNA-mediated cleavage of complementary mRNAs by RISC (PubMed:16289642, PubMed:17507929, PubMed:22791714). In cooperation with FMR1, regulates miRNA-mediated translational repression by AGO2 (PubMed:25464849). Restricts retrotransposition of long interspersed element-1 (LINE-1) in cooperation with TUT4 and TUT7 counteracting the RNA chaperone activity of L1RE1 (PubMed:23093941, PubMed:30122351). Facilitates LINE-1 uridylation by TUT4 and TUT7 (PubMed:30122351). Required for embryonic viability and for normal central nervous system development and function. Plays two critical roles in early brain development: suppresses retroelements in the nucleus by directly inhibiting cDNA synthesis, while regulates cytoskeletal mRNAs to influence neurite outgrowth in the cytosol (By similarity). May function as a messenger ribonucleoprotein (mRNP) clearance factor (PubMed:24726324).

Cellular Location

Cytoplasm, P-body. Cytoplasm, Cytoplasmic ribonucleoprotein granule. Cytoplasm, Stress granule. Nucleus {ECO:0000250|UniProtKB:P23249} Cytoplasm {ECO:0000250|UniProtKB:P23249}. Note=Co-enriched in cytoplasmic foci with TUT4 (PubMed:30122351). In developing neurons, localizes both in nucleus and cytoplasm, but in the adulthood it is only cytoplasmic (By similarity). After infection, relocalizes to the DENV replication complex in perinuclear regions (PubMed:27974568) {ECO:0000250|UniProtKB:P23249, ECO:0000269|PubMed:27974568, ECO:0000269|PubMed:30122351}

MOV10 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MOV10 Rabbit mAb - Images



