

**MOV10 Rabbit mAb**  
Catalog # AP76996**Specification****MOV10 Rabbit mAb - Product Information**

Application	WB
Primary Accession	<a href="#">Q9HCE1</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	113671

**MOV10 Rabbit mAb - Additional Information**

Gene ID 4343

**Other Names**

MOV10

**Dilution**

WB~~1/500-1/1000

**Format**

Liquid

**MOV10 Rabbit mAb - Protein Information**Name MOV10 ([HGNC:7200](#))

Synonyms KIAA1631

**Function**

5' to 3' RNA helicase that is involved in a number of cellular roles ranging from mRNA metabolism and translation, modulation of viral infectivity, inhibition of retrotransposition, or regulation of synaptic transmission (PubMed:[23093941](http://www.uniprot.org/citations/23093941)). Plays an important role in innate antiviral immunity by promoting type I interferon production (PubMed:[27016603](http://www.uniprot.org/citations/27016603), PubMed:[27974568](http://www.uniprot.org/citations/27974568), PubMed:[35157734](http://www.uniprot.org/citations/35157734)). Mechanistically, specifically uses IKKepsilon/IKBKE as the mediator kinase for IRF3 activation (PubMed:[27016603](http://www.uniprot.org/citations/27016603), PubMed:[35157734](http://www.uniprot.org/citations/35157734)). Blocks HIV-1 virus replication at a post-entry step (PubMed:[20215113](http://www.uniprot.org/citations/20215113)). Counteracts HIV-1 Vif-mediated degradation of APOBEC3G through its helicase activity by interfering with the ubiquitin-proteasome pathway (PubMed:[29258557](http://www.uniprot.org/citations/29258557)). Inhibits also hepatitis B virus/HBV replication by interacting with

HBV RNA and thereby inhibiting the early step of viral reverse transcription (PubMed:<a href="http://www.uniprot.org/citations/31722967" target="\_blank">31722967</a>). Contributes to UPF1 mRNA target degradation by translocation along 3' UTRs (PubMed:<a href="http://www.uniprot.org/citations/24726324" target="\_blank">24726324</a>). Required for microRNA (miRNA)-mediated gene silencing by the RNA-induced silencing complex (RISC). Required for both miRNA-mediated translational repression and miRNA-mediated cleavage of complementary mRNAs by RISC (PubMed:<a href="http://www.uniprot.org/citations/16289642" target="\_blank">16289642</a>, PubMed:<a href="http://www.uniprot.org/citations/17507929" target="\_blank">17507929</a>, PubMed:<a href="http://www.uniprot.org/citations/22791714" target="\_blank">22791714</a>). In cooperation with FMR1, regulates miRNA-mediated translational repression by AGO2 (PubMed:<a href="http://www.uniprot.org/citations/25464849" target="\_blank">25464849</a>). Restricts retrotransposition of long interspersed element-1 (LINE-1) in cooperation with TUT4 and TUT7 counteracting the RNA chaperone activity of L1RE1 (PubMed:<a href="http://www.uniprot.org/citations/23093941" target="\_blank">23093941</a>, PubMed:<a href="http://www.uniprot.org/citations/30122351" target="\_blank">30122351</a>). Facilitates LINE-1 uridylation by TUT4 and TUT7 (PubMed:<a href="http://www.uniprot.org/citations/30122351" target="\_blank">30122351</a>). Required for embryonic viability and for normal central nervous system development and function. Plays two critical roles in early brain development: suppresses retroelements in the nucleus by directly inhibiting cDNA synthesis, while regulates cytoskeletal mRNAs to influence neurite outgrowth in the cytosol (By similarity). May function as a messenger ribonucleoprotein (mRNP) clearance factor (PubMed:<a href="http://www.uniprot.org/citations/24726324" target="\_blank">24726324</a>).

#### Cellular Location

Cytoplasm, P-body. Cytoplasm, Cytoplasmic ribonucleoprotein granule. Cytoplasm, Stress granule. Nucleus {ECO:0000250|UniProtKB:P23249} Cytoplasm {ECO:0000250|UniProtKB:P23249}. Note=Co-enriched in cytoplasmic foci with TUT4 (PubMed:30122351). In developing neurons, localizes both in nucleus and cytoplasm, but in the adulthood it is only cytoplasmic (By similarity). After infection, relocates to the DENV replication complex in perinuclear regions (PubMed:27974568) {ECO:0000250|UniProtKB:P23249, ECO:0000269|PubMed:27974568, ECO:0000269|PubMed:30122351}

#### MOV10 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### MOV10 Rabbit mAb - Images



