

### IGF2BP2 Rabbit mAb

**Catalog # AP77030** 

# **Specification**

### IGF2BP2 Rabbit mAb - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW

WB
O9Y6M1
Human, Mouse, Rat
Rabbit
Monoclonal Antibody
66121

### IGF2BP2 Rabbit mAb - Additional Information

Gene ID 10644

Other Names IGF2BP2

**Dilution** WB~~1/500-1/1000

Format Liquid

## IGF2BP2 Rabbit mAb - Protein Information

Name IGF2BP2

Synonyms IMP2, VICKZ2

### **Function**

RNA-binding factor that recruits target transcripts to cytoplasmic protein-RNA complexes (mRNPs). This transcript 'caging' into mRNPs allows mRNA transport and transient storage. It also modulates the rate and location at which target transcripts encounter the translational apparatus and shields them from endonuclease attacks or microRNA-mediated degradation (By similarity). Preferentially binds to N6-methyladenosine (m6A)-containing mRNAs and increases their stability (PubMed:<a href="http://www.uniprot.org/citations/29476152" target="\_blank">29476152</a>). Binds to the 5'-UTR of the insulin-like growth factor 2 (IGF2) mRNAs (PubMed:<a href="http://www.uniprot.org/citations/9891060" target="\_blank">9891060</a>). Binding is isoform- specific. Binds to beta-actin/ACTB and MYC transcripts. Increases MYC mRNA stability by

isoform- specific. Binds to beta-actin/ACTB and MYC transcripts. Increases MYC mRNA stability by binding to the coding region instability determinant (CRD) and binding is enhanced by m6A-modification of the CRD (PubMed:<a href="http://www.uniprot.org/citations/29476152" target="\_blank">29476152</a>).

# **Cellular Location**

Nucleus. Cytoplasm. Cytoplasm, P-body. Cytoplasm, Stress granule. Note=Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Localizes at the connecting piece and the tail of





Tel: 858.875.1900 Fax: 858.875.1999

the spermatozoa. In response to cellular stress, such as oxidative stress, recruited to stress granules

## **Tissue Location**

Expressed in oocytes, granulosa cells of small and growing follicles, Leydig cells, spermatogonia and semen (at protein level). Expressed in testicular cancer (at protein level). Expressed weakly in heart, placenta, skeletal muscle, bone marrow, colon, kidney, salivary glands, testis and pancreas. Detected in fetal liver, fetal ovary, gonocytes and interstitial cells of the testis

## IGF2BP2 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# IGF2BP2 Rabbit mAb - Images

