

12 Lipoxygenase Rabbit mAb
Catalog # AP77032**Specification**

12 Lipoxygenase Rabbit mAb - Product Information

Application	WB
Primary Accession	P18054
Reactivity	Human, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	75694

12 Lipoxygenase Rabbit mAb - Additional Information

Gene ID 239

Other Names

ALOX12

Dilution

WB~~1/500-1/1000

Format

Liquid

12 Lipoxygenase Rabbit mAb - Protein InformationName ALOX12 ([HGNC:429](#))

Synonyms 12LO, LOG12

Function

Catalyzes the regio and stereo-specific incorporation of molecular oxygen into free and esterified polyunsaturated fatty acids generating lipid hydroperoxides that can be further reduced to the corresponding hydroxy species (PubMed: [17493578](http://www.uniprot.org/citations/17493578), PubMed: [18311922](http://www.uniprot.org/citations/18311922), PubMed: [1851637](http://www.uniprot.org/citations/1851637), PubMed: [32404334](http://www.uniprot.org/citations/32404334), PubMed: [8319693](http://www.uniprot.org/citations/8319693), PubMed: [8500694](http://www.uniprot.org/citations/8500694)). Mainly converts arachidonate ((5Z,8Z,11Z,14Z)-eicosatetraenoate) to the specific bioactive lipid (12S)-hydroperoxyeicosatetraenoate/(12S)-HPETE (PubMed: [17493578](http://www.uniprot.org/citations/17493578), PubMed: [22984144](http://www.uniprot.org/citations/22984144), PubMed: [24282679](http://www.uniprot.org/citations/24282679), PubMed: [8319693](http://www.uniprot.org/citations/8319693), PubMed: [8500694](http://www.uniprot.org/citations/8500694)). Through the

production of bioactive lipids like (12S)- HPETE it regulates different biological processes including platelet activation (PubMed:8319693, PubMed:8500694). It can also catalyze the epoxidation of double bonds of polyunsaturated fatty acids such as (14S)-hydroperoxy-docosahexaenoate/(14S)-HPDHA resulting in the formation of (13S,14S)-epoxy-DHA (PubMed:23504711). Furthermore, it may participate in the sequential oxidations of DHA ((4Z,7Z,10Z,13Z,16Z,19Z)-docosahexaenoate) to generate specialized pro- resolving mediators (SPMs) like resolvin D5 ((7S,17S)-diHPDHA) and (7S,14S)-diHPDHA, that actively down-regulate the immune response and have anti-aggregation properties with platelets (PubMed:32404334). An additional function involves a multistep process by which it transforms leukotriene A4/LTA4 into the bioactive lipids lipoxin A4/LXA4 and lipoxin B4/LXB4, both are vasoactive and LXA4 may regulate neutrophil function via occupancy of specific recognition sites (PubMed:8250832). Can also peroxidize linoleate ((9Z,12Z)-octadecadienoate) to (13S)- hydroperoxyoctadecadienoate/ (13S-HPODE) (By similarity). Due to its role in regulating both the expression of the vascular endothelial growth factor (VEGF, an angiogenic factor involved in the survival and metastasis of solid tumors) and the expression of integrin beta-1 (known to affect tumor cell migration and proliferation), it can be regarded as protumorigenic (PubMed:16638750, PubMed:22237009, PubMed:9751607). Important for cell survival, as it may play a role not only in proliferation but also in the prevention of apoptosis in vascular smooth muscle cells (PubMed:23578768).

Cellular Location

Cytoplasm, cytosol. Membrane. Note=Membrane association is stimulated by EGF

Tissue Location

Expressed in vascular smooth muscle cells.

12 Lipoxygenase Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

12 Lipoxygenase Rabbit mAb - Images



