

**Granulin Rabbit mAb**  
Catalog # AP77045**Specification**

---

**Granulin Rabbit mAb - Product Information**

|                   |                            |
|-------------------|----------------------------|
| Application       | <b>WB</b>                  |
| Primary Accession | <a href="#">P28799</a>     |
| Reactivity        | <b>Human</b>               |
| Host              | <b>Rabbit</b>              |
| Clonality         | <b>Monoclonal Antibody</b> |
| Calculated MW     | <b>63544</b>               |

**Granulin Rabbit mAb - Additional Information****Gene ID** 2896**Other Names**

GRN

**Dilution**

WB~~1/500-1/1000

**Format**

Liquid

**Granulin Rabbit mAb - Protein Information****Name** GRN ([HGNC:4601](#))**Function**

Secreted protein that acts as a key regulator of lysosomal function and as a growth factor involved in inflammation, wound healing and cell proliferation (PubMed:[12526812](http://www.uniprot.org/citations/12526812), PubMed:[18378771](http://www.uniprot.org/citations/18378771), PubMed:[28073925](http://www.uniprot.org/citations/28073925), PubMed:[28453791](http://www.uniprot.org/citations/28453791), PubMed:[28541286](http://www.uniprot.org/citations/28541286)). Regulates protein trafficking to lysosomes and, also the activity of lysosomal enzymes (PubMed:[28453791](http://www.uniprot.org/citations/28453791), PubMed:[28541286](http://www.uniprot.org/citations/28541286)). Facilitates also the acidification of lysosomes, causing degradation of mature CTSD by CTSB (PubMed:[28073925](http://www.uniprot.org/citations/28073925)). In addition, functions as a wound-related growth factor that acts directly on dermal fibroblasts and endothelial cells to promote division, migration and the formation of capillary-like tubule structures (By similarity). Also promotes epithelial cell proliferation by blocking TNF-mediated neutrophil activation preventing release of oxidants and proteases (PubMed:[12526812](http://www.uniprot.org/citations/12526812)). Moreover, modulates inflammation in neurons by preserving neurons survival, axonal outgrowth and

neuronal integrity (PubMed:<a href="http://www.uniprot.org/citations/18378771" target="\_blank">18378771</a>).

#### Cellular Location

Secreted. Lysosome Note=Endocytosed by SORT1 and delivered to lysosomes (PubMed:21092856, PubMed:28073925). Targeted to lysosome by PSAP via M6PR and LRP1, in both biosynthetic and endocytic pathways (PubMed:26370502, PubMed:28073925). Co-localized with GBA1 in the intracellular trafficking compartments until to lysosome (By similarity) {ECO:0000250|UniProtKB:P28798, ECO:0000269|PubMed:21092856, ECO:0000269|PubMed:26370502, ECO:0000269|PubMed:28073925}

#### Tissue Location

In myelogenous leukemic cell lines of promonocytic, promyelocytic, and proerythroid lineage, in fibroblasts, and very strongly in epithelial cell lines. Present in inflammatory cells and bone marrow. Highest levels in kidney

### Granulin Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Granulin Rabbit mAb - Images

