

Cullin 2 Rabbit mAb
Catalog # AP77098**Specification****Cullin 2 Rabbit mAb - Product Information**

Application	WB
Primary Accession	Q13617
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	86983

Cullin 2 Rabbit mAb - Additional Information**Gene ID** 8453**Other Names**
CUL2**Dilution**
WB~~1/500-1/1000**Format**
Liquid**Cullin 2 Rabbit mAb - Protein Information****Name** CUL2 ([HGNC:2552](#))**Function**

Core component of multiple cullin-RING-based ECS (ElonginB/C- CUL2/5-SOCS-box protein) E3 ubiquitin-protein ligase complexes, which mediate the ubiquitination of target proteins (PubMed:11384984, PubMed:26138980, PubMed:29775578, PubMed:29779948, PubMed:38326650). CUL2 serves as a rigid scaffold in the complex and may contribute to catalysis through positioning of the substrate and the E2 ubiquitin- conjugating enzyme (PubMed:10973499, PubMed:11384984, PubMed:12609982, PubMed:24076655, PubMed:9122164, PubMed:38326650). The E3 ubiquitin- protein ligase activity of the complex is dependent on the neddylation of the cullin subunit and is inhibited by the association of the deneddylated cullin subunit with TIP120A/CAND1 (PubMed:12609982, PubMed:11384984, PubMed:26138980, PubMed:29775578, PubMed:29779948, PubMed:38326650).

PubMed:24076655, PubMed:27565346, PubMed:38326650). The functional specificity of the ECS complex depends on the substrate recognition component (PubMed:10973499, PubMed:26138980, PubMed:29775578, PubMed:29779948, PubMed:9122164, PubMed:38326650). ECS(VHL) mediates the ubiquitination of hypoxia-inducible factor (HIF) (PubMed:10973499, PubMed:9122164). A number of ECS complexes (containing either KLHDC2, KLHDC3, KLHDC10, APPBP2, FEM1A, FEM1B or FEM1C as substrate-recognition component) are part of the DesCEND (destruction via C-end degrons) pathway, which recognizes a C-degron located at the extreme C terminus of target proteins, leading to their ubiquitination and degradation (PubMed:26138980, PubMed:29775578, PubMed:29779948). ECS complexes and ARIH1 collaborate in tandem to mediate ubiquitination of target proteins (PubMed:27565346). ECS(LRR1) ubiquitinates MCM7 and promotes CMG replisome disassembly by VCP and chromatin extraction during S- phase (By similarity).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q9D4H8}.

Cullin 2 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Cullin 2 Rabbit mAb - Images



