

**Presenilin 1 Rabbit mAb**  
Catalog # AP77146**Specification****Presenilin 1 Rabbit mAb - Product Information**

Application	WB
Primary Accession	<a href="#">P49768</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	52668

**Presenilin 1 Rabbit mAb - Additional Information**

Gene ID 5663

**Other Names**

PSEN1

**Dilution**

WB~~1/500-1/1000

**Format**

Liquid

**Presenilin 1 Rabbit mAb - Protein Information**

Name PSEN1

Synonyms AD3, PS1, PSNL1

**Function**

Catalytic subunit of the gamma-secretase complex, an endoprotease complex that catalyzes the intramembrane cleavage of integral membrane proteins such as Notch receptors and APP (amyloid- beta precursor protein) (PubMed: [10206644](http://www.uniprot.org/citations/10206644) target="\_blank">10206644</a>, PubMed: [10545183](http://www.uniprot.org/citations/10545183) target="\_blank">10545183</a>, PubMed: [10593990](http://www.uniprot.org/citations/10593990) target="\_blank">10593990</a>, PubMed: [10811883](http://www.uniprot.org/citations/10811883) target="\_blank">10811883</a>, PubMed: [10899933](http://www.uniprot.org/citations/10899933) target="\_blank">10899933</a>, PubMed: [12679784](http://www.uniprot.org/citations/12679784) target="\_blank">12679784</a>, PubMed: [12740439](http://www.uniprot.org/citations/12740439) target="\_blank">12740439</a>, PubMed: [15274632](http://www.uniprot.org/citations/15274632) target="\_blank">15274632</a>, PubMed: [20460383](http://www.uniprot.org/citations/20460383) target="\_blank">20460383</a>, PubMed: [25043039](http://www.uniprot.org/citations/25043039) target="\_blank">25043039</a>, PubMed: [26280335](http://www.uniprot.org/citations/26280335) target="\_blank">26280335</a>, PubMed: [28269784](http://www.uniprot.org/citations/28269784) target="\_blank">28269784</a>, PubMed: [30598546](http://www.uniprot.org/citations/30598546) target="\_blank">30598546</a>, PubMed: [30630874](http://www.uniprot.org/citations/30630874) target="\_blank">30630874</a>

target="\_blank">30630874</a>). Requires the presence of the other members of the gamma-secretase complex for protease activity (PubMed:<a href="http://www.uniprot.org/citations/15274632" target="\_blank">15274632</a>, PubMed:<a href="http://www.uniprot.org/citations/25043039" target="\_blank">25043039</a>, PubMed:<a href="http://www.uniprot.org/citations/26280335" target="\_blank">26280335</a>, PubMed:<a href="http://www.uniprot.org/citations/30598546" target="\_blank">30598546</a>, PubMed:<a href="http://www.uniprot.org/citations/30630874" target="\_blank">30630874</a>). Plays a role in Notch and Wnt signaling cascades and regulation of downstream processes via its role in processing key regulatory proteins, and by regulating cytosolic CTNNB1 levels (PubMed:<a href="http://www.uniprot.org/citations/10593990" target="\_blank">10593990</a>, PubMed:<a href="http://www.uniprot.org/citations/10811883" target="\_blank">10811883</a>, PubMed:<a href="http://www.uniprot.org/citations/10899933" target="\_blank">10899933</a>, PubMed:<a href="http://www.uniprot.org/citations/9738936" target="\_blank">9738936</a>). Stimulates cell-cell adhesion via its interaction with CDH1; this stabilizes the complexes between CDH1 (E-cadherin) and its interaction partners CTNNB1 (beta-catenin), CTNND1 and JUP (gamma-catenin) (PubMed:<a href="http://www.uniprot.org/citations/11953314" target="\_blank">11953314</a>). Under conditions of apoptosis or calcium influx, cleaves CDH1 (PubMed:<a href="http://www.uniprot.org/citations/11953314" target="\_blank">11953314</a>). This promotes the disassembly of the complexes between CDH1 and CTNND1, JUP and CTNNB1, increases the pool of cytoplasmic CTNNB1, and thereby negatively regulates Wnt signaling (PubMed:<a href="http://www.uniprot.org/citations/11953314" target="\_blank">11953314</a>, PubMed:<a href="http://www.uniprot.org/citations/9738936" target="\_blank">9738936</a>). Required for normal embryonic brain and skeleton development, and for normal angiogenesis (By similarity). Mediates the proteolytic cleavage of EphB2/CTF1 into EphB2/CTF2 (PubMed:<a href="http://www.uniprot.org/citations/17428795" target="\_blank">17428795</a>, PubMed:<a href="http://www.uniprot.org/citations/28269784" target="\_blank">28269784</a>). The holoprotein functions as a calcium-leak channel that allows the passive movement of calcium from endoplasmic reticulum to cytosol and is therefore involved in calcium homeostasis (PubMed:<a href="http://www.uniprot.org/citations/16959576" target="\_blank">16959576</a>, PubMed:<a href="http://www.uniprot.org/citations/25394380" target="\_blank">25394380</a>). Involved in the regulation of neurite outgrowth (PubMed:<a href="http://www.uniprot.org/citations/15004326" target="\_blank">15004326</a>, PubMed:<a href="http://www.uniprot.org/citations/20460383" target="\_blank">20460383</a>). Is a regulator of presynaptic facilitation, spike transmission and synaptic vesicles replenishment in a process that depends on gamma-secretase activity. It acts through the control of SYT7 presynaptic expression (By similarity).

### Cellular Location

Endoplasmic reticulum. Endoplasmic reticulum membrane; Multi-pass membrane protein. Golgi apparatus membrane; Multi-pass membrane protein. Cytoplasmic granule. Cell membrane; Multi-pass membrane protein. Cell projection, growth cone. Early endosome. Early endosome membrane; Multi-pass membrane protein. Cell projection, neuron projection. Cell projection, axon {ECO:0000250|UniProtKB:Q4JIM4}. Synapse {ECO:0000250|UniProtKB:Q4JIM4}. Note=Translocates with bound NOTCH1 from the endoplasmic reticulum and/or Golgi to the cell surface (PubMed:10593990). Colocalizes with CDH1/2 at sites of cell-cell contact. Colocalizes with CTNNB1 in the endoplasmic reticulum and the proximity of the plasma membrane (PubMed:9738936). Also present in azurophil granules of neutrophils (PubMed:11987239). Colocalizes with UBQLN1 in the cell membrane and in cytoplasmic juxtannuclear structures called aggresomes (PubMed:21143716).

### Tissue Location

Detected in azurophilic granules in neutrophils and in platelet cytoplasmic granules (at protein level) (PubMed:11987239) Expressed in a wide range of tissues including various regions of the brain, liver, spleen and lymph nodes (PubMed:7596406, PubMed:8574969, PubMed:8641442).

## Presenilin 1 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Presenilin 1 Rabbit mAb - Images

