

# IRAK1 Rabbit mAb

**Catalog # AP77196** 

# **Specification**

#### IRAK1 Rabbit mAb - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW

WB
P51617
Human
Rabbit
Monoclonal Antibody

76537

### IRAK1 Rabbit mAb - Additional Information

**Gene ID 3654** 

Other Names IRAK1

**Dilution** WB~~1/500-1/1000

Format Liquid

## **IRAK1** Rabbit mAb - Protein Information

Name IRAK1 (HGNC:6112)


**Synonyms IRAK** 

## **Function**

Serine/threonine-protein kinase that plays a critical role in initiating innate immune response against foreign pathogens. Involved in Toll-like receptor (TLR) and IL-1R signaling pathways. Is rapidly recruited by MYD88 to the receptor-signaling complex upon TLR activation. Association with MYD88 leads to IRAK1 phosphorylation by IRAK4 and subsequent autophosphorylation and kinase activation. Phosphorylates E3 ubiquitin ligases Pellino proteins (PELI1, PELI2 and PELI3) to promote pellino-mediated polyubiquitination of IRAK1. Then, the ubiquitin-binding domain of IKBKG/NEMO binds to polyubiquitinated IRAK1 bringing together the IRAK1-MAP3K7/TAK1-TRAF6 complex and the NEMO-IKKA-IKKB complex. In turn, MAP3K7/TAK1 activates IKKs (CHUK/IKKA and IKBKB/IKKB) leading to NF-kappa-B nuclear translocation and activation. Alternatively, phosphorylates TIRAP to promote its ubiquitination and subsequent degradation. Phosphorylates the interferon regulatory factor 7 (IRF7) to induce its activation and translocation to the nucleus, resulting in transcriptional activation of type I IFN genes, which drive the cell in an antiviral state. When sumoylated, translocates to the nucleus and phosphorylates STAT3.

## **Cellular Location**

Cytoplasm. Nucleus. Lipid droplet Note=Translocates to the nucleus when sumoylated.





Tel: 858.875.1900 Fax: 858.875.1999

RSAD2/viperin recruits it to the lipid droplet (By similarity).

## **Tissue Location**

Isoform 1 and isoform 2 are ubiquitously expressed in all tissues examined, with isoform 1 being more strongly expressed than isoform 2.

## **IRAK1** Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# IRAK1 Rabbit mAb - Images

