

**IL-10 Rabbit mAb**  
Catalog # AP77246**Specification**

---

**IL-10 Rabbit mAb - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">P22301</a>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Monoclonal Antibody</b>
Calculated MW	<b>20517</b>

**IL-10 Rabbit mAb - Additional Information****Gene ID** 3586**Other Names**

IL10

**Dilution**

WB~~1/500-1/1000

**Format**

Liquid

**IL-10 Rabbit mAb - Protein Information****Name** IL10**Function**

Major immune regulatory cytokine that acts on many cells of the immune system where it has profound anti-inflammatory functions, limiting excessive tissue disruption caused by inflammation. Mechanistically, IL10 binds to its heterotetrameric receptor comprising IL10RA and IL10RB leading to JAK1 and STAT2-mediated phosphorylation of STAT3 (PubMed:<a href="http://www.uniprot.org/citations/16982608" target="\_blank">16982608</a>). In turn, STAT3 translocates to the nucleus where it drives expression of anti-inflammatory mediators (PubMed:<a href="http://www.uniprot.org/citations/18025162" target="\_blank">18025162</a>). Targets antigen-presenting cells (APCs) such as macrophages and monocytes and inhibits their release of pro- inflammatory cytokines including granulocyte-macrophage colony- stimulating factor /GM-CSF, granulocyte colony-stimulating factor/G- CSF, IL-1 alpha, IL-1 beta, IL-6, IL-8 and TNF-alpha (PubMed:<a href="http://www.uniprot.org/citations/11564774" target="\_blank">11564774</a>, PubMed:<a href="http://www.uniprot.org/citations/1940799" target="\_blank">1940799</a>, PubMed:<a href="http://www.uniprot.org/citations/7512027" target="\_blank">7512027</a>). Interferes also with antigen presentation by reducing the expression of MHC-class II and co- stimulatory molecules, thereby inhibiting their ability to induce T cell activation (PubMed:<a href="http://www.uniprot.org/citations/8144879" target="\_blank">8144879</a>). In addition, controls the inflammatory response of macrophages by reprogramming essential metabolic pathways including mTOR signaling (By similarity).

**Cellular Location**

Secreted.

**Tissue Location**

Produced by a variety of cell lines, including T- cells, macrophages, mast cells and other cell types

**IL-10 Rabbit mAb - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**IL-10 Rabbit mAb - Images**