

14-3-3 sigma Rabbit mAb
Catalog # AP77287**Specification**

14-3-3 sigma Rabbit mAb - Product Information

Application	WB
Primary Accession	P31947
Reactivity	Human, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	27774

14-3-3 sigma Rabbit mAb - Additional Information

Gene ID 2810

Other Names

SFN

Dilution

WB~~1/500-1/1000

Format

Liquid

14-3-3 sigma Rabbit mAb - Protein Information

Name SFN

Synonyms HME1 {ECO:0000303|PubMed:1390337}

Function

Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways (PubMed: [15731107](http://www.uniprot.org/citations/15731107), PubMed: [22634725](http://www.uniprot.org/citations/22634725), PubMed: [28202711](http://www.uniprot.org/citations/28202711), PubMed: [37797010](http://www.uniprot.org/citations/37797010)). Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif (PubMed: [15731107](http://www.uniprot.org/citations/15731107), PubMed: [22634725](http://www.uniprot.org/citations/22634725), PubMed: [28202711](http://www.uniprot.org/citations/28202711), PubMed: [37797010](http://www.uniprot.org/citations/37797010)). Binding generally results in the modulation of the activity of the binding partner (PubMed: [15731107](http://www.uniprot.org/citations/15731107), PubMed: [22634725](http://www.uniprot.org/citations/22634725), PubMed: [28202711](http://www.uniprot.org/citations/28202711), PubMed: [37797010](http://www.uniprot.org/citations/37797010)). Promotes

cytosolic retention of GBP1 GTPase by binding to phosphorylated GBP1, thereby inhibiting the innate immune response (PubMed:37797010). Also acts as a TP53/p53-regulated inhibitor of G2/M progression (PubMed:9659898). When bound to KRT17, regulates protein synthesis and epithelial cell growth by stimulating Akt/mTOR pathway (By similarity). Acts to maintain desmosome cell junction adhesion in epithelial cells via interacting with and sequestering PKP3 to the cytoplasm, thereby restricting its translocation to existing desmosome structures and therefore maintaining desmosome protein homeostasis (PubMed:24124604). Also acts to facilitate PKP3 exchange at desmosome plaques, thereby maintaining keratinocyte intercellular adhesion (PubMed:29678907). May also regulate MDM2 autoubiquitination and degradation and thereby activate p53/TP53 (PubMed:18382127).

Cellular Location

Cytoplasm. Nucleus {ECO:0000250|UniProtKB:O70456} Secreted. Note=May be secreted by a non- classical secretory pathway.

Tissue Location

Present mainly in tissues enriched in stratified squamous keratinizing epithelium.

14-3-3 sigma Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

14-3-3 sigma Rabbit mAb - Images



