

Rab11A Rabbit mAb
Catalog # AP77443**Specification**

Rab11A Rabbit mAb - Product Information

Application	WB
Primary Accession	P62491
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	24394

Rab11A Rabbit mAb - Additional Information**Gene ID** 8766**Other Names**
RAB11A**Dilution**
WB~~1/500-1/1000**Format**
Liquid**Rab11A Rabbit mAb - Protein Information****Name** RAB11A ([HGNC:9760](#))**Function**

The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different set of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion (PubMed: [15601896](http://www.uniprot.org/citations/15601896), PubMed: [15689490](http://www.uniprot.org/citations/15689490), PubMed: [17462998](http://www.uniprot.org/citations/17462998), PubMed: [19542231](http://www.uniprot.org/citations/19542231), PubMed: [20026645](http://www.uniprot.org/citations/20026645), PubMed: [20890297](http://www.uniprot.org/citations/20890297), PubMed: [21282656](http://www.uniprot.org/citations/21282656)). The small Rab GTPase RAB11A regulates endocytic recycling (PubMed: [20026645](http://www.uniprot.org/citations/20026645)). Forms a functional Rab11/FIP3/dynein complex that regulates the movement of peripheral sorting endosomes (SE) along microtubule tracks toward the microtubule organizing center/centrosome, generating the endosomal recycling compartment (ERC) (PubMed: [20026645](http://www.uniprot.org/citations/20026645)). Acts as a major regulator of membrane delivery during cytokinesis (PubMed: [20026645](http://www.uniprot.org/citations/20026645)).

href="http://www.uniprot.org/citations/15601896" target="_blank">15601896). Together with MYO5B and RAB8A participates in epithelial cell polarization. Together with RAB3IP, RAB8A, the exocyst complex, PARD3, PRKCI, ANXA2, CDC42 and DNMBP promotes transcytosis of PODXL to the apical membrane initiation sites (AMIS), apical surface formation and lumenogenesis. Together with MYO5B participates in CFTR trafficking to the plasma membrane and TF (Transferrin) recycling in nonpolarized cells. Required in a complex with MYO5B and RAB11FIP2 for the transport of NPC1L1 to the plasma membrane. Participates in the sorting and basolateral transport of CDH1 from the Golgi apparatus to the plasma membrane. Regulates the recycling of FCGRT (receptor of Fc region of monomeric Ig G) to basolateral membranes. May also play a role in melanosome transport and release from melanocytes (PubMed:15689490, PubMed:17462998, PubMed:19542231, PubMed:20890297, PubMed:21282656). Promotes Rabin8/RAB3IP preciliary vesicular trafficking to mother centriole by forming a ciliary targeting complex containing Rab11, ASAP1, Rabin8/RAB3IP, RAB11FIP3 and ARF4, thereby regulating ciliogenesis initiation (PubMed:25673879, PubMed:31204173). On the contrary, upon LPAR1 receptor signaling pathway activation, interaction with phosphorylated WDR44 prevents Rab11-RAB3IP-RAB11FIP3 complex formation and cilia growth (PubMed:31204173). Participates in the export of a subset of neosynthesized proteins through a Rab8-Rab10-Rab11- endosomal dependent export route via interaction with WDR44 (PubMed:32344433).

Cellular Location

Cell membrane; Lipid-anchor. Endosome membrane. Recycling endosome membrane; Lipid-anchor. Cleavage furrow. Cytoplasmic vesicle, phagosome. Cytoplasmic vesicle membrane. Golgi apparatus. Golgi apparatus, trans-Golgi network. Note=Localized to WDR44-positive endosomes and tubules (PubMed:32344433). Translocates with RAB11FIP2 from the vesicles of the endocytic recycling compartment (ERC) to the plasma membrane (PubMed:11994279). During interphase, localized in vesicles continuously moving from peripheral sorting endosomes towards the pericentrosomal ERC (PubMed:20026645). Localizes to the cleavage furrow (PubMed:15601896). Colocalizes with PARD3, PRKCI, EXOC5, OCLN, PODXL and RAB8A in apical membrane initiation sites (AMIS) during the generation of apical surface and lumenogenesis (PubMed:20890297) Recruited to phagosomes containing S.aureus or M.tuberculosis (PubMed:21255211). Localized to rhodopsin transport carriers when interacting with RAB11AFIP3 and ASAP1 in photoreceptors (PubMed:25673879).

Rab11A Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Rab11A Rabbit mAb - Images

