

**ARHE Antibody (Center)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP7751c**

**Specification**

---

**ARHE Antibody (Center) - Product Information**

Application	<b>WB, IHC-P,E</b>
Primary Accession	<a href="#">P61587</a>
Other Accession	<a href="#">Q6SA80</a> , <a href="#">O77683</a> , <a href="#">P61588</a>
Reactivity	<b>Human, Mouse, Rat</b>
Predicted	<b>Pig</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit IgG</b>
Antigen Region	<b>133-165</b>

**ARHE Antibody (Center) - Additional Information**

**Gene ID** 390

**Other Names**

Rho-related GTP-binding protein RhoE, Protein MemB, Rho family GTPase 3, Rho-related GTP-binding protein Rho8, Rnd3, RND3, ARHE, RHO8, RHOE

**Target/Specificity**

This ARHE antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 133-165 amino acids from the Central region of human ARHE.

**Dilution**

WB~~1:2000  
IHC-P~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

ARHE Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**ARHE Antibody (Center) - Protein Information**

**Name** RND3

**Synonyms** ARHE, RHO8, RHOE

**Function** Binds GTP but lacks intrinsic GTPase activity and is resistant to Rho-specific GTPase-activating proteins.

**Cellular Location**

Golgi apparatus membrane; Peripheral membrane protein

**Tissue Location**


Ubiquitous.

**ARHE Antibody (Center) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**ARHE Antibody (Center) - Images**



Western blot analysis of ARHE Antibody (Center)(Cat. 3AP7751c) in mouse brain tissue lysates (35ug/lane). ARHE (arrow) was detected using the purified Pab.



All lanes : Anti-ARHE Antibody (Center) at 1:2000 dilution Lane 1: mouse brain lysate Lane 2: rat brain lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 27 kDa Blocking/Dilution buffer: 5% NFD/MTBST.



Formalin-fixed and paraffin-embedded human hepatocarcinoma tissue reacted with ARHE antibody (Center) (Cat.#AP7751c), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

### **ARHE Antibody (Center) - Background**

Members of the Rho family of Ras-related GTPases, such as ARHE, regulate the organization of the actin cytoskeleton in response to extracellular growth factors. Like Ras (MIM 190020), Rho family members appear to cycle between an inactive GDP-bound form and an active GTP-bound form. Three major regulators of Rho activity have been identified: RhoGDIs, which interact with the GDP-bound Rho proteins to keep them in a resting complex (see MIM 601925); GEFs, which promote GDP/GTP exchange leading to activation of Rho proteins (see MIM 601855); and GAPs, which stimulate GTP hydrolysis and return the activated Rho protein to its inactive form (see MIM 602680) (Nobes et al., 1998 [PubMed 9531558]).

### **ARHE Antibody (Center) - References**

- Pinner,S., Nat. Cell Biol. 10 (2), 127-137 (2008)
- Poch,E., Exp. Cell Res. 313 (4), 719-731 (2007)
- Ongusaha,P.P., Curr. Biol. 16 (24), 2466-2472 (2006)