

# CD3D Rabbit mAb

Catalog # AP77585

### Specification

## CD3D Rabbit mAb - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW

WB, IHC, IF <u>P04234</u> Human Rabbit Monoclonal Antibody 18930

### CD3D Rabbit mAb - Additional Information

Gene ID 915

Other Names CD3D

**Dilution** WB~~1/500-1/1000 IHC~~1/50-1/100 IF~~1/50-1/200

Format Liquid

## CD3D Rabbit mAb - Protein Information

Name CD3D

Synonyms T3D

#### Function

Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in adaptive immune response. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR- mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain. Upon TCR engagement, these motifs become phosphorylated by Src family protein tyrosine kinases LCK and FYN, resulting in the activation of downstream signaling pathways (PubMed:<a href="http://www.uniprot.org/citations/2470098"">http://www.uniprot.org/citations/2470098</a>

target="\_blank">2470098</a>). In addition of this role of signal transduction in T-cell activation, CD3D plays an essential role in thymocyte differentiation. Indeed, participates in correct intracellular TCR-CD3 complex assembly and surface expression. In absence of a functional TCR-CD3 complex, thymocytes are unable to differentiate properly. Interacts with CD4 and CD8 and thus serves to establish a functional link between the TCR and coreceptors CD4 and CD8, which is needed for activation and positive selection of CD4 or CD8 T-cells (PubMed:<a href="http://www.uniprot.org/citations/12215456" target="\_blank">12215456</a>).



#### **Cellular Location**

Cell membrane; Single-pass type I membrane protein

**Tissue Location** 

CD3D is mostly present on T-lymphocytes with its TCR-CD3 partners. Present also in fetal NK-cells

### **CD3D Rabbit mAb - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

### CD3D Rabbit mAb - Images







