

**CD19 Rabbit mAb**  
Catalog # AP77609**Specification**

---

**CD19 Rabbit mAb - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">P15391</a>
Reactivity	<b>Human, Mouse, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Monoclonal Antibody</b>
Calculated MW	<b>61128</b>

**CD19 Rabbit mAb - Additional Information****Gene ID** 930**Other Names**

CD19

**Dilution**

WB~~1/500-1/1000

**Format**

Liquid

**CD19 Rabbit mAb - Protein Information****Name** CD19**Function**

Functions as a coreceptor for the B-cell antigen receptor complex (BCR) on B-lymphocytes (PubMed: [29523808](http://www.uniprot.org/citations/29523808)). Decreases the threshold for activation of downstream signaling pathways and for triggering B-cell responses to antigens (PubMed: [1373518](http://www.uniprot.org/citations/1373518), PubMed: [16672701](http://www.uniprot.org/citations/16672701), PubMed: [2463100](http://www.uniprot.org/citations/2463100)). Activates signaling pathways that lead to the activation of phosphatidylinositol 3-kinase and the mobilization of intracellular Ca(2+) stores (PubMed: [12387743](http://www.uniprot.org/citations/12387743), PubMed: [16672701](http://www.uniprot.org/citations/16672701), PubMed: [9317126](http://www.uniprot.org/citations/9317126), PubMed: [9382888](http://www.uniprot.org/citations/9382888)). Is not required for early steps during B cell differentiation in the blood marrow (PubMed: [9317126](http://www.uniprot.org/citations/9317126)). Required for normal differentiation of B-1 cells (By similarity). Required for normal B cell differentiation and proliferation in response to antigen challenges (PubMed: [1373518](http://www.uniprot.org/citations/1373518), PubMed: [2463100](http://www.uniprot.org/citations/2463100)). Required for

normal levels of serum immunoglobulins, and for production of high-affinity antibodies in response to antigen challenge (PubMed:<a href="http://www.uniprot.org/citations/12387743" target="\_blank">12387743</a>, PubMed:<a href="http://www.uniprot.org/citations/16672701" target="\_blank">16672701</a>, PubMed:<a href="http://www.uniprot.org/citations/9317126" target="\_blank">9317126</a>).

#### Cellular Location

Cell membrane; Single-pass type I membrane protein. Membrane raft  
{ECO:0000250|UniProtKB:P25918}; Single-pass type I membrane protein  
{ECO:0000250|UniProtKB:P25918}

#### Tissue Location

Detected on marginal zone and germinal center B cells in lymph nodes (PubMed:2463100).  
Detected on blood B cells (at protein level) (PubMed:16672701, PubMed:2463100)

### CD19 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### CD19 Rabbit mAb - Images

