

**BMP2 Rabbit mAb**  
Catalog # AP77621**Specification**

---

**BMP2 Rabbit mAb - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">P12643</a>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Monoclonal Antibody</b>
Calculated MW	<b>44702</b>

**BMP2 Rabbit mAb - Additional Information**

Gene ID 650

**Other Names**

BMP2

**Dilution**

WB~~1/500-1/1000

**Format**

Liquid

**BMP2 Rabbit mAb - Protein Information**

Name BMP2

Synonyms BMP2A

**Function**

Growth factor of the TGF-beta superfamily that plays essential roles in many developmental processes, including cardiogenesis, neurogenesis, and osteogenesis (PubMed: [18436533](http://www.uniprot.org/citations/18436533), PubMed: [24362451](http://www.uniprot.org/citations/24362451), PubMed: [31019025](http://www.uniprot.org/citations/31019025)). Induces cartilage and bone formation (PubMed: [3201241](http://www.uniprot.org/citations/3201241)). Initiates the canonical BMP signaling cascade by associating with type I receptor BMPR1A and type II receptor BMPR2 (PubMed: [15064755](http://www.uniprot.org/citations/15064755), PubMed: [17295905](http://www.uniprot.org/citations/17295905), PubMed: [18436533](http://www.uniprot.org/citations/18436533)). Once all three components are bound together in a complex at the cell surface, BMPR2 phosphorylates and activates BMPR1A (PubMed: [7791754](http://www.uniprot.org/citations/7791754)). In turn, BMPR1A propagates signal by phosphorylating SMAD1/5/8 that travel to the nucleus and act as activators and repressors of transcription of target genes. Also acts to promote expression of HAMP, via the interaction with its receptor

BMPR1A/ALK3 (PubMed:<a href="http://www.uniprot.org/citations/31800957" target="\_blank">31800957</a>). Can also signal through non-canonical pathways such as ERK/MAP kinase signaling cascade that regulates osteoblast differentiation (PubMed:<a href="http://www.uniprot.org/citations/16771708" target="\_blank">16771708</a>, PubMed:<a href="http://www.uniprot.org/citations/20851880" target="\_blank">20851880</a>). Also stimulates the differentiation of myoblasts into osteoblasts via the EIF2AK3-EIF2A-ATF4 pathway by stimulating EIF2A phosphorylation which leads to increased expression of ATF4 which plays a central role in osteoblast differentiation (PubMed:<a href="http://www.uniprot.org/citations/24362451" target="\_blank">24362451</a>). Acts as a positive regulator of odontoblast differentiation during mesenchymal tooth germ formation, expression is repressed during the bell stage by MSX1-mediated inhibition of CTNNB1 signaling (By similarity).

#### Cellular Location

Secreted.

#### Tissue Location

Particularly abundant in lung, spleen and colon and in low but significant levels in heart, brain, placenta, liver, skeletal muscle, kidney, pancreas, prostate, ovary and small intestine

#### BMP2 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### BMP2 Rabbit mAb - Images

