

NeuroD1 Rabbit mAb

Catalog # AP77658

Specification

NeuroD1 Rabbit mAb - Product Information

Application Primary Accession Reactivity Host Clonality WB
O13562
Human, Mouse, Rat
Rabbit
Monoclonal Antibody
39920

NeuroD1 Rabbit mAb - Additional Information

Gene ID 4760

Calculated MW

Other Names NEUROD1

Dilution WB~~1/500-1/1000

Format Liquid

NeuroD1 Rabbit mAb - Protein Information

Name NEUROD1

Synonyms BHLHA3, NEUROD

Function

Acts as a transcriptional activator: mediates transcriptional activation by binding to E box-containing promoter consensus core sequences 5'-CANNTG-3'. Associates with the p300/CBP transcription coactivator complex to stimulate transcription of the secretin gene as well as the gene encoding the cyclin-dependent kinase inhibitor CDKN1A. Contributes to the regulation of several cell differentiation pathways, like those that promote the formation of early retinal ganglion cells, inner ear sensory neurons, granule cells forming either the cerebellum or the dentate gyrus cell layer of the hippocampus, endocrine islet cells of the pancreas and enteroendocrine cells of the small intestine. Together with PAX6 or SIX3, is required for the regulation of amacrine cell fate specification. Also required for dendrite morphogenesis and maintenance in the cerebellar cortex. Associates with chromatin to enhancer regulatory elements in genes encoding key transcriptional regulators of neurogenesis (By similarity).

Cellular Location

Cytoplasm. Nucleus {ECO:0000255|PROSITE-ProRule:PRU00981, ECO:0000269|PubMed:14752053} Note=In pancreatic islet cells, shuttles to the nucleus in response to glucose stimulation (By similarity). Colocalizes with NR0B2 in the nucleus.



NeuroD1 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

NeuroD1 Rabbit mAb - Images

