

**PRAS40 Rabbit mAb**  
Catalog # AP77762**Specification****PRAS40 Rabbit mAb - Product Information**

|                   |                            |
|-------------------|----------------------------|
| Application       | <b>WB</b>                  |
| Primary Accession | <a href="#">Q96B36</a>     |
| Reactivity        | <b>Human, Mouse, Rat</b>   |
| Host              | <b>Rabbit</b>              |
| Clonality         | <b>Monoclonal Antibody</b> |
| Calculated MW     | <b>27383</b>               |

**PRAS40 Rabbit mAb - Additional Information****Gene ID** 84335**Other Names**

AKT1S1

**Dilution**

WB~~1/500-1/1000

**Format**

Liquid

**PRAS40 Rabbit mAb - Protein Information****Name** AKT1S1 {ECO:0000312|EMBL:AAH16043.1}**Function**

Negative regulator of the mechanistic target of rapamycin complex 1 (mTORC1), an evolutionarily conserved central nutrient sensor that stimulates anabolic reactions and macromolecule biosynthesis to promote cellular biomass generation and growth (PubMed:<a href="http://www.uniprot.org/citations/17277771" target="\_blank">17277771</a>, PubMed:<a href="http://www.uniprot.org/citations/17386266" target="\_blank">17386266</a>, PubMed:<a href="http://www.uniprot.org/citations/17510057" target="\_blank">17510057</a>, PubMed:<a href="http://www.uniprot.org/citations/29236692" target="\_blank">29236692</a>). In absence of insulin and nutrients, AKT1S1 associates with the mTORC1 complex and directly inhibits mTORC1 activity by blocking the MTOR substrate- recruitment site (PubMed:<a href="http://www.uniprot.org/citations/29236692" target="\_blank">29236692</a>). In response to insulin and nutrients, AKT1S1 dissociates from mTORC1 (PubMed:<a href="http://www.uniprot.org/citations/17386266" target="\_blank">17386266</a>, PubMed:<a href="http://www.uniprot.org/citations/18372248" target="\_blank">18372248</a>). Its activity is dependent on its phosphorylation state and binding to 14-3-3 (PubMed:<a href="http://www.uniprot.org/citations/16174443" target="\_blank">16174443</a>, PubMed:<a href="http://www.uniprot.org/citations/18372248" target="\_blank">18372248</a>). May also play a role in nerve growth factor-mediated neuroprotection (By similarity).

**Cellular Location**

Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q9D1F4}. Note=Found in the cytosolic fraction of the brain. {ECO:0000250|UniProtKB:Q9D1F4}

**Tissue Location**

Widely expressed with highest levels of expression in liver and heart. Expressed at higher levels in cancer cell lines (e.g. A-549 and HeLa) than in normal cell lines (e.g. HEK293)

**PRAS40 Rabbit mAb - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**PRAS40 Rabbit mAb - Images**