

CD33 Rabbit mAb
Catalog # AP77800**Specification**

CD33 Rabbit mAb - Product Information

Application	WB
Primary Accession	P20138
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	39825

CD33 Rabbit mAb - Additional Information**Gene ID** 945**Other Names**
CD33**Dilution**
WB~~1/500-1/1000**Format**
Liquid**CD33 Rabbit mAb - Protein Information****Name** CD33**Synonyms** SIGLEC3**Function**

Sialic-acid-binding immunoglobulin-like lectin (Siglec) that plays a role in mediating cell-cell interactions and in maintaining immune cells in a resting state (PubMed: [10611343](http://www.uniprot.org/citations/10611343), PubMed: [11320212](http://www.uniprot.org/citations/11320212), PubMed: [15597323](http://www.uniprot.org/citations/15597323)). Preferentially recognizes and binds alpha-2,3- and more avidly alpha-2,6-linked sialic acid-bearing glycans (PubMed: [7718872](http://www.uniprot.org/citations/7718872)). Upon engagement of ligands such as C1q or sialylated glycoproteins, two immunoreceptor tyrosine-based inhibitory motifs (ITIMs) located in CD33 cytoplasmic tail are phosphorylated by Src-like kinases such as LCK (PubMed: [10887109](http://www.uniprot.org/citations/10887109), PubMed: [28325905](http://www.uniprot.org/citations/28325905)). These phosphorylations provide docking sites for the recruitment and activation of protein-tyrosine phosphatases PTPN6/SHP-1 and PTPN11/SHP-2 (PubMed: [10206955](http://www.uniprot.org/citations/10206955), PubMed: [10556798](http://www.uniprot.org/citations/10556798), PubMed: [10887109](http://www.uniprot.org/citations/10887109)).

In turn, these phosphatases regulate downstream pathways through dephosphorylation of signaling molecules (PubMed:10206955, PubMed:10887109). One of the repressive effect of CD33 on monocyte activation requires phosphoinositide 3-kinase/PI3K (PubMed:15597323).

Cellular Location

[Isoform CD33M]: Cell membrane; Single-pass type I membrane protein

Tissue Location

Monocytic/myeloid lineage cells. In the brain, CD33 is mainly expressed on microglial cells

CD33 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD33 Rabbit mAb - Images

