

Transcription Factor E3 Rabbit mAb
Catalog # AP77851**Specification****Transcription Factor E3 Rabbit mAb - Product Information**

Application	IHC
Primary Accession	P19532
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	61521

Transcription Factor E3 Rabbit mAb - Additional Information

Gene ID 7030

Other Names

TFE3

Dilution

IHC~~1/50-1/100

Format

Liquid

Transcription Factor E3 Rabbit mAb - Protein Information

Name TFE3 {ECO:0000303|PubMed:9393982, ECO:0000312|HGNC:HGNC:11752}

Function

Transcription factor that acts as a master regulator of lysosomal biogenesis and immune response (PubMed: [2338243](http://www.uniprot.org/citations/2338243), PubMed: [24448649](http://www.uniprot.org/citations/24448649), PubMed: [29146937](http://www.uniprot.org/citations/29146937), PubMed: [30733432](http://www.uniprot.org/citations/30733432), PubMed: [31672913](http://www.uniprot.org/citations/31672913), PubMed: [37079666](http://www.uniprot.org/citations/37079666)). Specifically recognizes and binds E-box sequences (5'-CANNTG-3'); efficient DNA-binding requires dimerization with itself or with another MiT/TFE family member such as TFEB or MITF (PubMed: [24448649](http://www.uniprot.org/citations/24448649)). Involved in the cellular response to amino acid availability by acting downstream of MTOR: in the presence of nutrients, TFE3 phosphorylation by MTOR promotes its inactivation (PubMed: [24448649](http://www.uniprot.org/citations/24448649), PubMed: [31672913](http://www.uniprot.org/citations/31672913), PubMed: [36608670](http://www.uniprot.org/citations/36608670)). Upon starvation or lysosomal stress, inhibition of MTOR induces TFE3 dephosphorylation, resulting in transcription factor activity (PubMed: [24448649](http://www.uniprot.org/citations/24448649), PubMed: [31672913](http://www.uniprot.org/citations/31672913)).

target="_blank">31672913, PubMed:36608670). Specifically recognizes and binds the CLEAR-box sequence (5'-GTCACGTGAC-3') present in the regulatory region of many lysosomal genes, leading to activate their expression, thereby playing a central role in expression of lysosomal genes (PubMed:24448649). Maintains the pluripotent state of embryonic stem cells by promoting the expression of genes such as ESRRB; mTOR- dependent TFE3 cytosolic retention and inactivation promotes exit from pluripotency (By similarity). Required to maintain the naive pluripotent state of hematopoietic stem cell; mTOR-dependent cytoplasmic retention of TFE3 promotes the exit of hematopoietic stem cell from pluripotency (PubMed:30733432). TFE3 activity is also involved in the inhibition of neuronal progenitor differentiation (By similarity). Acts as a positive regulator of browning of adipose tissue by promoting expression of target genes; mTOR-dependent phosphorylation promotes cytoplasmic retention of TFE3 and inhibits browning of adipose tissue (By similarity). In association with TFEB, activates the expression of CD40L in T-cells, thereby playing a role in T-cell- dependent antibody responses in activated CD4(+) T-cells and thymus- dependent humoral immunity (By similarity). Specifically recognizes the MUE3 box, a subset of E-boxes, present in the immunoglobulin enhancer (PubMed:2338243). It also binds very well to a USF/MLTF site (PubMed:2338243). Promotes TGF-beta-induced transcription of COL1A2; via its interaction with TSC22D1 at E-boxes in the gene proximal promoter (By similarity). May regulate lysosomal positioning in response to nutrient deprivation by promoting the expression of PIP4P1 (PubMed:29146937).

Cellular Location

Cytoplasm, cytosol. Nucleus. Lysosome membrane. Note=When nutrients are present, recruited to the lysosomal membrane via association with GDP-bound RagC/RRAGC (or RagD/RRAGD): it is then phosphorylated by MTOR (PubMed:24448649, PubMed:37079666). Phosphorylation by MTOR prevents nuclear translocation and promotes ubiquitination and degradation (PubMed:22692423, PubMed:30733432, PubMed:36608670, PubMed:37079666) Conversely, inhibition of mTORC1, starvation and lysosomal disruption, promotes dephosphorylation and translocation to the nucleus (PubMed:22692423, PubMed:30733432, PubMed:37079666)

Tissue Location

Ubiquitous in fetal and adult tissues.

Transcription Factor E3 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Transcription Factor E3 Rabbit mAb - Images



