

MAD2L1 Rabbit mAb
Catalog # AP77922**Specification****MAD2L1 Rabbit mAb - Product Information**

Application	WB
Primary Accession	Q13257
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	23510

MAD2L1 Rabbit mAb - Additional Information

Gene ID 4085

Other Names
MAD2L1**Dilution**
WB~~1/500-1/1000**Format**
Liquid**MAD2L1 Rabbit mAb - Protein Information****Name** MAD2L1**Synonyms** MAD2**Function**

Component of the spindle-assembly checkpoint that prevents the onset of anaphase until all chromosomes are properly aligned at the metaphase plate (PubMed: [15024386](http://www.uniprot.org/citations/15024386), PubMed: [29162720](http://www.uniprot.org/citations/29162720)). In the closed conformation (C-MAD2) forms a heterotetrameric complex with MAD1L1 at unattached kinetochores during prometaphase, the complex recruits open conformation molecules of MAD2L1 (O-MAD2) and then promotes the conversion of O-MAD2 to C-MAD2 (PubMed: [29162720](http://www.uniprot.org/citations/29162720)). Required for the execution of the mitotic checkpoint which monitors the process of kinetochore-spindle attachment and inhibits the activity of the anaphase promoting complex by sequestering CDC20 until all chromosomes are aligned at the metaphase plate (PubMed: [10700282](http://www.uniprot.org/citations/10700282), PubMed: [11804586](http://www.uniprot.org/citations/11804586), PubMed: [15024386](http://www.uniprot.org/citations/15024386)).

Cellular Location

Nucleus. Chromosome, centromere, kinetochore. Cytoplasm. Cytoplasm, cytoskeleton, spindle pole Note=Recruited by MAD1L1 to unattached kinetochores (Probable) Recruited to the nuclear pore complex by TPR during interphase Recruited to kinetochores in late prometaphase after BUB1, CENPF, BUB1B and CENPE. Kinetochore association requires the presence of NEK2 Kinetochore association is repressed by UBD. Sequestered to the cytoplasm upon interaction with isoform 3 of MAD1L1 (PubMed:19010891) {ECO:0000269|PubMed:19010891, ECO:0000305}

MAD2L1 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MAD2L1 Rabbit mAb - Images

