

Sprouty 4 Rabbit mAb
Catalog # AP77958**Specification**

Sprouty 4 Rabbit mAb - Product Information

Application	WB
Primary Accession	O9C004
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	32541

Sprouty 4 Rabbit mAb - Additional Information**Gene ID** 81848**Other Names**
SPRY4**Dilution**
WB~~1/500-1/1000**Format**
Liquid**Sprouty 4 Rabbit mAb - Protein Information****Name** SPRY4**Function**

Suppresses the insulin receptor and EGFR-transduced MAPK signaling pathway, but does not inhibit MAPK activation by a constitutively active mutant Ras (PubMed:[12027893](http://www.uniprot.org/citations/12027893)). Probably impairs the formation of GTP-Ras (PubMed:[12027893](http://www.uniprot.org/citations/12027893)). Inhibits Ras-independent, but not Ras-dependent, activation of RAF1 (PubMed:[12717443](http://www.uniprot.org/citations/12717443)). Represses integrin-mediated cell spreading via inhibition of TESK1-mediated phosphorylation of cofilin (PubMed:[15584898](http://www.uniprot.org/citations/15584898)).

Cellular Location

Cytoplasm. Cell projection, ruffle membrane; Peripheral membrane protein; Cytoplasmic side. Note=Found in the cytoplasm in unstimulated cells but is translocated to the membrane ruffles in cells stimulated with EGF (epidermal growth factor) (By similarity). Colocalizes with TESK1 in vesicular spots in the cytoplasm (PubMed:15584898) {ECO:0000250|UniProtKB:Q9WTP2, ECO:0000269|PubMed:15584898}

Sprouty 4 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Sprouty 4 Rabbit mAb - Images

