

HIF1AN Rabbit mAb
Catalog # AP78027**Specification**

HIF1AN Rabbit mAb - Product Information

Application	WB
Primary Accession	Q9NWT6
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	40285

HIF1AN Rabbit mAb - Additional Information

Gene ID 55662

Other Names
HIF1AN**Dilution**
WB~~1/500-1/1000**Format**
Liquid**HIF1AN Rabbit mAb - Protein Information****Name** HIF1AN**Synonyms** FIH1**Function**

Hydroxylates HIF-1 alpha at 'Asn-803' in the C-terminal transactivation domain (CAD). Functions as an oxygen sensor and, under normoxic conditions, the hydroxylation prevents interaction of HIF-1 with transcriptional coactivators including Cbp/p300-interacting transactivator. Involved in transcriptional repression through interaction with HIF1A, VHL and histone deacetylases. Hydroxylates specific Asn residues within ankyrin repeat domains (ARD) of NFKB1, NFKBIA, NOTCH1, ASB4, PPP1R12A and several other ARD-containing proteins. Also hydroxylates Asp and His residues within ARDs of ANK1 and TNKS2, respectively. Negatively regulates NOTCH1 activity, accelerating myogenic differentiation. Positively regulates ASB4 activity, promoting vascular differentiation.

Cellular Location

Nucleus. Cytoplasm. Cytoplasm, perinuclear region. Note=Mainly cytoplasmic localization, but interaction with NOTCH1 results in nuclear localization and interaction with ABPA3 results in perinuclear localization in macrophages

HIF1AN Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HIF1AN Rabbit mAb - Images

