

**Cytochrome P450 17A1 Rabbit mAb**  
Catalog # AP78069**Specification****Cytochrome P450 17A1 Rabbit mAb - Product Information**

Application	WB
Primary Accession	<a href="#">P05093</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	57371

**Cytochrome P450 17A1 Rabbit mAb - Additional Information**

Gene ID 1586

**Other Names**  
CYP17A1**Dilution**  
WB~~1/500-1/1000**Format**  
Liquid**Cytochrome P450 17A1 Rabbit mAb - Protein Information****Name** CYP17A1 {ECO:0000303|PubMed:19793597, ECO:0000312|HGNC:HGNC:2593}**Function**

A cytochrome P450 monooxygenase involved in corticoid and androgen biosynthesis (PubMed: [22266943](http://www.uniprot.org/citations/22266943), PubMed: [25301938](http://www.uniprot.org/citations/25301938), PubMed: [27339894](http://www.uniprot.org/citations/27339894), PubMed: [9452426](http://www.uniprot.org/citations/9452426)). Catalyzes 17-alpha hydroxylation of C21 steroids, which is common for both pathways. A second oxidative step, required only for androgen synthesis, involves an acyl-carbon cleavage. The 17-alpha hydroxy intermediates, as part of adrenal glucocorticoids biosynthesis pathway, are precursors of cortisol (Probable) (PubMed: [25301938](http://www.uniprot.org/citations/25301938), PubMed: [9452426](http://www.uniprot.org/citations/9452426)). Hydroxylates steroid hormones, pregnenolone and progesterone to form 17-alpha hydroxy metabolites, followed by the cleavage of the C17-C20 bond to form C19 steroids, dehydroepiandrosterone (DHEA) and androstenedione (PubMed: [22266943](http://www.uniprot.org/citations/22266943), PubMed: [25301938](http://www.uniprot.org/citations/25301938), PubMed: [27339894](http://www.uniprot.org/citations/27339894), PubMed: [36640554](http://www.uniprot.org/citations/36640554), PubMed: [9452426](http://www.uniprot.org/citations/9452426)). Has 16-alpha

hydroxylase activity. Catalyzes 16-alpha hydroxylation of 17-alpha hydroxy pregnenolone, followed by the cleavage of the C17-C20 bond to form 16-alpha-hydroxy DHEA (PubMed:<a href="http://www.uniprot.org/citations/36640554" target="\_blank">36640554</a>). Also 16-alpha hydroxylates androgens, relevant for estriol synthesis (PubMed:<a href="http://www.uniprot.org/citations/25301938" target="\_blank">25301938</a>, PubMed:<a href="http://www.uniprot.org/citations/27339894" target="\_blank">27339894</a>). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (CPR; NADPH-ferrihemoprotein reductase) (PubMed:<a href="http://www.uniprot.org/citations/22266943" target="\_blank">22266943</a>, PubMed:<a href="http://www.uniprot.org/citations/25301938" target="\_blank">25301938</a>, PubMed:<a href="http://www.uniprot.org/citations/27339894" target="\_blank">27339894</a>, PubMed:<a href="http://www.uniprot.org/citations/9452426" target="\_blank">9452426</a>).

#### Cellular Location

Endoplasmic reticulum membrane. Microsome membrane

#### Cytochrome P450 17A1 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Cytochrome P450 17A1 Rabbit mAb - Images

