

**CD16 Rabbit mAb**  
Catalog # AP78175**Specification**

---

**CD16 Rabbit mAb - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">P08637</a>
Reactivity	<b>Human, Mouse, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Monoclonal Antibody</b>
Calculated MW	<b>29089</b>

**CD16 Rabbit mAb - Additional Information****Gene ID** 2214**Other Names**  
FCGR3A**Dilution**  
WB~~1/500-1/1000**Format**  
Liquid**CD16 Rabbit mAb - Protein Information****Name** FCGR3A {ECO:0000303|PubMed:23006327}**Function**

Receptor for the invariable Fc fragment of immunoglobulin gamma (IgG). Optimally activated upon binding of clustered antigen-IgG complexes displayed on cell surfaces, triggers lysis of antibody-coated cells, a process known as antibody-dependent cellular cytotoxicity (ADCC). Does not bind free monomeric IgG, thus avoiding inappropriate effector cell activation in the absence of antigenic trigger (PubMed: [11711607](http://www.uniprot.org/citations/11711607), PubMed: [21768335](http://www.uniprot.org/citations/21768335), PubMed: [22023369](http://www.uniprot.org/citations/22023369), PubMed: [24412922](http://www.uniprot.org/citations/24412922), PubMed: [25786175](http://www.uniprot.org/citations/25786175), PubMed: [25816339](http://www.uniprot.org/citations/25816339), PubMed: [28652325](http://www.uniprot.org/citations/28652325), PubMed: [8609432](http://www.uniprot.org/citations/8609432), PubMed: [9242542](http://www.uniprot.org/citations/9242542)). Mediates IgG effector functions on natural killer (NK) cells. Binds antigen-IgG complexes generated upon infection and triggers NK cell-dependent cytokine production and degranulation to limit viral load and propagation. Involved in the generation of memory- like adaptive NK cells capable to produce high amounts of IFNG and to efficiently eliminate virus-infected cells via ADCC (PubMed: [23006327](#))

href="http://www.uniprot.org/citations/24412922" target="\_blank">24412922</a>, PubMed:<a href="http://www.uniprot.org/citations/25786175" target="\_blank">25786175</a>). Regulates NK cell survival and proliferation, in particular by preventing NK cell progenitor apoptosis (PubMed:<a href="http://www.uniprot.org/citations/29967280" target="\_blank">29967280</a>, PubMed:<a href="http://www.uniprot.org/citations/9916693" target="\_blank">9916693</a>). Fc-binding subunit that associates with CD247 and/or FCER1G adapters to form functional signaling complexes. Following the engagement of antigen-IgG complexes, triggers phosphorylation of immunoreceptor tyrosine-based activation motif (ITAM)-containing adapters with subsequent activation of phosphatidylinositol 3-kinase signaling and sustained elevation of intracellular calcium that ultimately drive NK cell activation. The ITAM-dependent signaling coupled to receptor phosphorylation by PKC mediates robust intracellular calcium flux that leads to production of pro-inflammatory cytokines, whereas in the absence of receptor phosphorylation it mainly activates phosphatidylinositol 3-kinase signaling leading to cell degranulation (PubMed:<a href="http://www.uniprot.org/citations/1825220" target="\_blank">1825220</a>, PubMed:<a href="http://www.uniprot.org/citations/23024279" target="\_blank">23024279</a>, PubMed:<a href="http://www.uniprot.org/citations/2532305" target="\_blank">2532305</a>). Costimulates NK cells and trigger lysis of target cells independently of IgG binding (PubMed:<a href="http://www.uniprot.org/citations/10318937" target="\_blank">10318937</a>, PubMed:<a href="http://www.uniprot.org/citations/23006327" target="\_blank">23006327</a>). Mediates the antitumor activities of therapeutic antibodies. Upon ligation on monocytes triggers TNFA-dependent ADCC of IgG-coated tumor cells (PubMed:<a href="http://www.uniprot.org/citations/27670158" target="\_blank">27670158</a>). Mediates enhanced ADCC in response to afucosylated IgGs (PubMed:<a href="http://www.uniprot.org/citations/34485821" target="\_blank">34485821</a>).

#### **Cellular Location**

Cell membrane; Single-pass type I membrane protein. Secreted. Note=Exists also as a soluble receptor

#### **Tissue Location**

Expressed in natural killer cells (at protein level) (PubMed:2526846). Expressed in a subset of circulating monocytes (at protein level) (PubMed:27670158).

### **CD16 Rabbit mAb - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **CD16 Rabbit mAb - Images**



