

**G3BP Rabbit mAb**  
Catalog # AP78234**Specification**

---

**G3BP Rabbit mAb - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">O13283</a>
Reactivity	<b>Human, Mouse, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Monoclonal Antibody</b>
Calculated MW	<b>52164</b>

**G3BP Rabbit mAb - Additional Information**

Gene ID 10146

**Other Names**  
G3BP1**Dilution**  
WB~~1/500-1/1000**Format**  
Liquid**G3BP Rabbit mAb - Protein Information****Name** G3BP1 {ECO:0000303|PubMed:23279204, ECO:0000312|HGNC:HGNC:30292}**Function**

Protein involved in various processes, such as stress granule formation and innate immunity (PubMed: <a href="http://www.uniprot.org/citations/12642610" target="\_blank">12642610</a>, PubMed: <a href="http://www.uniprot.org/citations/20180778" target="\_blank">20180778</a>, PubMed: <a href="http://www.uniprot.org/citations/23279204" target="\_blank">23279204</a>, PubMed: <a href="http://www.uniprot.org/citations/30510222" target="\_blank">30510222</a>, PubMed: <a href="http://www.uniprot.org/citations/30804210" target="\_blank">30804210</a>). Plays an essential role in stress granule formation (PubMed: <a href="http://www.uniprot.org/citations/12642610" target="\_blank">12642610</a>, PubMed: <a href="http://www.uniprot.org/citations/20180778" target="\_blank">20180778</a>, PubMed: <a href="http://www.uniprot.org/citations/23279204" target="\_blank">23279204</a>, PubMed: <a href="http://www.uniprot.org/citations/32302570" target="\_blank">32302570</a>, PubMed: <a href="http://www.uniprot.org/citations/32302571" target="\_blank">32302571</a>, PubMed: <a href="http://www.uniprot.org/citations/32302572" target="\_blank">32302572</a>, PubMed: <a href="http://www.uniprot.org/citations/34739333" target="\_blank">34739333</a>, PubMed: <a href="http://www.uniprot.org/citations/35977029" target="\_blank">35977029</a>, PubMed: <a href="http://www.uniprot.org/citations/36183834" target="\_blank">36183834</a>, PubMed: <a href="http://www.uniprot.org/citations/36279435" target="\_blank">36279435</a>, PubMed: <a href="http://www.uniprot.org/citations/36692217" target="\_blank">36692217</a>, PubMed: <a href="http://www.uniprot.org/citations/12642610" target="\_blank">12642610</a>, PubMed: <a href="http://www.uniprot.org/citations/20180778" target="\_blank">20180778</a>, PubMed: <a href="http://www.uniprot.org/citations/23279204" target="\_blank">23279204</a>, PubMed: <a href="http://www.uniprot.org/citations/30510222" target="\_blank">30510222</a>, PubMed: <a href="http://www.uniprot.org/citations/30804210" target="\_blank">30804210</a>).

[37379838](http://www.uniprot.org/citations/37379838)). Stress granules are membraneless compartments that store mRNAs and proteins, such as stalled translation pre-initiation complexes, in response to stress (PubMed:[12642610](http://www.uniprot.org/citations/12642610), PubMed:[20180778](http://www.uniprot.org/citations/20180778), PubMed:[23279204](http://www.uniprot.org/citations/23279204), PubMed:[27022092](http://www.uniprot.org/citations/27022092), PubMed:[32302570](http://www.uniprot.org/citations/32302570), PubMed:[32302571](http://www.uniprot.org/citations/32302571), PubMed:[32302572](http://www.uniprot.org/citations/32302572), PubMed:[36279435](http://www.uniprot.org/citations/36279435), PubMed:[37379838](http://www.uniprot.org/citations/37379838)). Promotes formation of stress granules phase-separated membraneless compartment by undergoing liquid-liquid phase separation (LLPS) upon unfolded RNA-binding: functions as a molecular switch that triggers RNA-dependent LLPS in response to a rise in intracellular free RNA concentrations (PubMed:[32302570](http://www.uniprot.org/citations/32302570), PubMed:[32302571](http://www.uniprot.org/citations/32302571), PubMed:[32302572](http://www.uniprot.org/citations/32302572), PubMed:[34739333](http://www.uniprot.org/citations/34739333), PubMed:[36279435](http://www.uniprot.org/citations/36279435), PubMed:[36692217](http://www.uniprot.org/citations/36692217)). Also acts as an ATP- and magnesium-dependent helicase: unwinds DNA/DNA, RNA/DNA, and RNA/RNA substrates with comparable efficiency (PubMed:[9889278](http://www.uniprot.org/citations/9889278)). Acts unidirectionally by moving in the 5' to 3' direction along the bound single-stranded DNA (PubMed:[9889278](http://www.uniprot.org/citations/9889278)). Unwinds preferentially partial DNA and RNA duplexes having a 17 bp annealed portion and either a hanging 3' tail or hanging tails at both 5'- and 3'-ends (PubMed:[9889278](http://www.uniprot.org/citations/9889278)). Plays an essential role in innate immunity by promoting CGAS and RIGI activity (PubMed:[30510222](http://www.uniprot.org/citations/30510222), PubMed:[30804210](http://www.uniprot.org/citations/30804210)). Participates in the DNA-triggered cGAS/STING pathway by promoting the DNA binding and activation of CGAS (PubMed:[30510222](http://www.uniprot.org/citations/30510222)). Triggers the condensation of cGAS, a process probably linked to the formation of membrane-less organelles (PubMed:[34779554](http://www.uniprot.org/citations/34779554) target="\_blank">34779554</a>). Enhances also RIGI-induced type I interferon production probably by helping RIGI at sensing pathogenic RNA (PubMed:[30804210](http://www.uniprot.org/citations/30804210) target="\_blank">30804210</a>). May also act as a phosphorylation- dependent sequence-specific endoribonuclease in vitro: Cleaves exclusively between cytosine and adenine and cleaves MYC mRNA preferentially at the 3'-UTR (PubMed:[11604510](http://www.uniprot.org/citations/11604510) target="\_blank">11604510</a>).

### Cellular Location

Cytoplasm, cytosol. Perikaryon {ECO:0000250|UniProtKB:P97855}. Cytoplasm, Stress granule. Nucleus Note=Cytoplasmic in proliferating cells (PubMed:11604510). Cytosolic and partially nuclear in resting cells (PubMed:11604510). Recruited to stress granules in response to arsenite treatment (PubMed:12642610, PubMed:20180778). The unphosphorylated form is recruited to stress granules (PubMed:12642610). HRAS signaling contributes to this process by regulating G3BP dephosphorylation (PubMed:12642610)

### Tissue Location

Ubiquitous..

### G3BP Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### G3BP Rabbit mAb - Images

