

DDX39B Rabbit mAb
Catalog # AP78253**Specification****DDX39B Rabbit mAb - Product Information**

Application	WB
Primary Accession	Q13838
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	48991

DDX39B Rabbit mAb - Additional Information

Gene ID 7919

Other Names
DDX39B**Dilution**
WB~~1/500-1/1000**Format**
Liquid**DDX39B Rabbit mAb - Protein Information****Name** DDX39B ([HGNC:13917](#))**Synonyms** BAT1, UAP56**Function**

Involved in nuclear export of spliced and unspliced mRNA (PubMed:[15833825](http://www.uniprot.org/citations/15833825), PubMed:[15998806](http://www.uniprot.org/citations/15998806), PubMed:[17190602](http://www.uniprot.org/citations/17190602)). Component of the TREX complex which is thought to couple mRNA transcription, processing and nuclear export, and specifically associates with spliced mRNA and not with unspliced pre-mRNA (PubMed:[15833825](http://www.uniprot.org/citations/15833825), PubMed:[15998806](http://www.uniprot.org/citations/15998806), PubMed:[17190602](http://www.uniprot.org/citations/17190602)). The TREX complex is recruited to spliced mRNAs by a transcription-independent mechanism, binds to mRNA upstream of the exon-junction complex (EJC) and is recruited in a splicing- and cap-dependent manner to a region near the 5' end of the mRNA where it functions in mRNA export to the cytoplasm via the TAP/NXF1 pathway (PubMed:[15833825](http://www.uniprot.org/citations/15833825), PubMed:[15998806](http://www.uniprot.org/citations/15998806), PubMed:[17190602](http://www.uniprot.org/citations/17190602)). The

THOC1-THOC2- THOC3 core complex alone is sufficient to promote ATPase activity of DDX39B; in the complex THOC2 is the only component that directly interacts with DDX39B (PubMed:33191911). Associates with SARNP/CIP29, which facilitates RNA binding of DDX39B and likely plays a role in mRNA export (PubMed:37578863). May undergo several rounds of ATP hydrolysis during assembly of TREX to drive subsequent loading of components such as ALYREF/THOC4 and CHTOP onto mRNA. Also associates with pre-mRNA independent of ALYREF/THOC4. Involved in the nuclear export of intronless mRNA; the ATP-bound form is proposed to recruit export adapter ALYREF/THOC4 to intronless mRNA; its ATPase activity is cooperatively stimulated by RNA and ALYREF/THOC4 and ATP hydrolysis is thought to trigger the dissociation from RNA to allow the association of ALYREF/THOC4 and the NXF1-NXT1 heterodimer. Involved in transcription elongation and genome stability.

Cellular Location

Nucleus. Nucleus speckle. Cytoplasm. Note=Can translocate to the cytoplasm in the presence of MX1. TREX complex assembly seems to occur in regions surrounding nuclear speckles known as perispeckles

DDX39B Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DDX39B Rabbit mAb - Images

