

**DcR2 Rabbit mAb**  
Catalog # AP78306**Specification**

---

**DcR2 Rabbit mAb - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">O9UBN6</a>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Monoclonal Antibody</b>
Calculated MW	<b>41849</b>

**DcR2 Rabbit mAb - Additional Information****Gene ID** 8793**Other Names**  
TNFRSF10D**Dilution**  
WB~~1/500-1/1000**Format**  
Liquid**DcR2 Rabbit mAb - Protein Information****Name** TNFRSF10D ([HGNC:11907](#))**Function**

Receptor for the cytotoxic ligand TRAIL (PubMed: [9430226](http://www.uniprot.org/citations/9430226)). Contains a truncated death domain and hence is not capable of inducing apoptosis but protects against TRAIL-mediated apoptosis (PubMed: [9537512](http://www.uniprot.org/citations/9537512)). Reports are contradictory with regards to its ability to induce the NF-kappa-B pathway. According to PubMed: [9382840](http://www.uniprot.org/citations/9382840), it cannot but according to PubMed: [9430226](http://www.uniprot.org/citations/9430226), it can induce the NF-kappa-B pathway (PubMed: [9382840](http://www.uniprot.org/citations/9382840), PubMed: [9430226](http://www.uniprot.org/citations/9430226)).

**Cellular Location**

Membrane; Single-pass type I membrane protein

**Tissue Location**

Widely expressed, in particular in fetal kidney, lung and liver, and in adult testis and liver. Also expressed in peripheral blood leukocytes, colon and small intestine, ovary, prostate, thymus,

spleen, pancreas, kidney, lung, placenta and heart

### DcR2 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### DcR2 Rabbit mAb - Images

