

Uracil DNA Glycosylase Rabbit mAb
Catalog # AP78312

Specification

Uracil DNA Glycosylase Rabbit mAb - Product Information

Application	WB
Primary Accession	P13051
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	34645

Uracil DNA Glycosylase Rabbit mAb - Additional Information

Gene ID 7374

Other Names
UNG

Dilution
WB~~1/500-1/1000

Format
Liquid

Uracil DNA Glycosylase Rabbit mAb - Protein Information

Name UNG {ECO:0000255|HAMAP-Rule:MF_03166}

Function
Excises uracil residues from the DNA which can arise as a result of misincorporation of dUMP residues by DNA polymerase or due to deamination of cytosine.

Cellular Location
[Isoform 1]: Mitochondrion.

Tissue Location
Isoform 1 is widely expressed with the highest expression in skeletal muscle, heart and testicles. Isoform 2 has the highest expression levels in tissues containing proliferating cells

Uracil DNA Glycosylase Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Uracil DNA Glycosylase Rabbit mAb - Images

