

SARS Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7834B

Specification

SARS Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	P49591
Other Accession	Q6P799 , P26638 , Q4R4U9 , P26636 , Q9GMB8 , P13642
Reactivity	Human
Predicted	Bovine, Hamster, Monkey, Mouse, Rabbit, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	58777
Antigen Region	394-422

SARS Antibody (C-term) - Additional Information

Gene ID 6301

Other Names

Serine--tRNA ligase, cytoplasmic, Seryl-tRNA synthetase, SerRS, Seryl-tRNA(Ser/Sec) synthetase, SARS, SERS

Target/Specificity

This SARS antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 394-422 amino acids from the C-terminal region of human SARS.

Dilution

WB~~1:1000
IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SARS Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

SARS Antibody (C-term) - Protein Information

Name SARS1 ([HGNC:10537](#))

Synonyms SARS, SERS

Function Catalyzes the attachment of serine to tRNA(Ser) in a two-step reaction: serine is first activated by ATP to form Ser-AMP and then transferred to the acceptor end of tRNA(Ser) (PubMed:[22353712](#), PubMed:[24095058](#), PubMed:[26433229](#), PubMed:[28236339](#), PubMed:[34570399](#), PubMed:[36041817](#), PubMed:[9431993](#)). Is probably also able to aminoacylate tRNA(Sec) with serine, to form the misacylated tRNA L-seryl-tRNA(Sec), which will be further converted into selenocysteinyl-tRNA(Sec) (PubMed:[26433229](#), PubMed:[28236339](#), PubMed:[34570399](#), PubMed:[9431993](#)). In the nucleus, binds to the VEGFA core promoter and prevents MYC binding and transcriptional activation by MYC (PubMed:[24940000](#)). Recruits SIRT2 to the VEGFA promoter, promoting deacetylation of histone H4 at 'Lys- 16' (H4K16). Thereby, inhibits the production of VEGFA and sprouting angiogenesis mediated by VEGFA (PubMed:[19423847](#), PubMed:[19423848](#), PubMed:[24940000](#)).

Cellular Location

Cytoplasm. Nucleus Note=Predominantly cytoplasmic, but a minor proportion is also found in the nucleus.

Tissue Location


Brain..

SARS Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SARS Antibody (C-term) - Images



Western blot analysis of anti-SARS Antibody (C-term) (Cat.#AP7834b) in NCI-H460 cell line lysates (35ug/lane). SARS(arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human brain tissue reacted with SARS antibody (C-term) (Cat.#AP7834b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

SARS Antibody (C-term) - Background

Seryl-tRNA synthetase belongs to the class II amino-acyl tRNA family. This enzyme catalyzes the transfer of L-serine to tRNA (Ser) and is related to bacterial and yeast counterparts.

SARS Antibody (C-term) - References

- Shimada,N., J. Biol. Chem. 276 (50), 46770-46778 (2001)
- Shah,Z.H., Hum. Mutat. 17 (5), 433-434 (2001)
- Heckl,M., FEBS Lett. 427 (3), 315-319 (1998)