

**HLA F Rabbit mAb**  
Catalog # AP78428**Specification**

---

**HLA F Rabbit mAb - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">P30511</a>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Monoclonal Antibody</b>
Calculated MW	<b>39062</b>

**HLA F Rabbit mAb - Additional Information****Gene ID** 3134**Other Names**

HLA-F

**Dilution**

WB~~1/500-1/1000

**Format**

Liquid

**HLA F Rabbit mAb - Protein Information****Name** HLA-F**Function**

Non-classical major histocompatibility class Ib molecule postulated to play a role in immune surveillance, immune tolerance and inflammation. Functions in two forms, as a heterotrimeric complex with B2M/beta-2 microglobulin and a peptide (peptide-bound HLA-F-B2M) and as an open conformer (OC) devoid of peptide and B2M (peptide-free OC). In complex with B2M, presents non-canonical self-peptides carrying post-translational modifications, particularly phosphorylated self-peptides. Peptide-bound HLA-F-B2M acts as a ligand for LILRB1 inhibitory receptor, a major player in maternal-fetal tolerance. Peptide-free OC acts as a ligand for KIR3DS1 and KIR3DL2 receptors (PubMed: [28636952](http://www.uniprot.org/citations/28636952)). Upon interaction with activating KIR3DS1 receptor on NK cells, triggers NK cell degranulation and anti-viral cytokine production (PubMed: [27455421](http://www.uniprot.org/citations/27455421)). Through interaction with KIR3DL2 receptor, inhibits NK and T cell effector functions (PubMed: [24018270](http://www.uniprot.org/citations/24018270)). May interact with other MHC class I OCs to cross-present exogenous viral, tumor or minor histocompatibility antigens to cytotoxic CD8+ T cells, triggering effector and memory responses (PubMed: [23851683](http://www.uniprot.org/citations/23851683)). May play a role in inflammatory responses in the peripheral nervous system. Through interaction with KIR3DL2, may protect motor neurons from astrocyte-induced toxicity (PubMed: [23851683](#)).

<http://www.uniprot.org/citations/26928464> target="\_blank">26928464</a>).

### Cellular Location

Cell membrane; Single-pass type I membrane protein. Early endosome membrane. Lysosome membrane. Note=For cross-presentation transits from the cell surface through endosomal pathway to lysosomes, where the peptide is generated from internalized exogenous antigen

### Tissue Location

Expressed in resting B cells (at protein level). Expressed in secondary lymphoid organs rich in B and T cells such as the tonsils, spleen, and thymus (at protein level) (PubMed:10605026, PubMed:11169396). Expressed in the endothelial cells of the tonsils (PubMed:11169396). Expressed on activated lymphoid cells including B cells, NK cells, CD4+ T cells and memory T cells (at protein level) (PubMed:20865824, PubMed:27455421). Expressed in motor neurons of spinal cord (PubMed:26928464).

### HLA F Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### HLA F Rabbit mAb - Images

