

RAGE Rabbit mAb
Catalog # AP78520**Specification**

RAGE Rabbit mAb - Product Information

Application	WB
Primary Accession	Q15109
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	42803

RAGE Rabbit mAb - Additional Information**Gene ID** 177**Other Names**

AGER

Format

Liquid

RAGE Rabbit mAb - Protein Information**Name** AGER**Synonyms** RAGE**Function**

Cell surface pattern recognition receptor that senses endogenous stress signals with a broad ligand repertoire including advanced glycation end products, S100 proteins, high-mobility group box 1 protein/HMGB1, amyloid beta/APP oligomers, nucleic acids, phospholipids and glycosaminoglycans (PubMed: [27572515](http://www.uniprot.org/citations/27572515), PubMed: [28515150](http://www.uniprot.org/citations/28515150), PubMed: [34743181](http://www.uniprot.org/citations/34743181)). Advanced glycosylation end products are nonenzymatically glycosylated proteins which accumulate in vascular tissue in aging and at an accelerated rate in diabetes (PubMed: [21565706](http://www.uniprot.org/citations/21565706)). These ligands accumulate at inflammatory sites during the pathogenesis of various diseases, including diabetes, vascular complications, neurodegenerative disorders, and cancers and RAGE transduces their binding into pro-inflammatory responses. Upon ligand binding, uses TIRAP and MYD88 as adapters to transduce the signal ultimately leading to the induction of inflammatory cytokines IL6, IL8 and TNFalpha through activation of NF-kappa-B (PubMed: [21829704](http://www.uniprot.org/citations/21829704), PubMed: [33436632](http://www.uniprot.org/citations/33436632)). Interaction with S100A12 on endothelium, mononuclear phagocytes, and lymphocytes triggers cellular activation, with generation of key pro-inflammatory mediators (PubMed: [19386136](http://www.uniprot.org/citations/19386136)). Interaction

with S100B after myocardial infarction may play a role in myocyte apoptosis by activating ERK1/2 and p53/TP53 signaling (By similarity). Contributes to the translocation of amyloid- beta peptide (ABPP) across the cell membrane from the extracellular to the intracellular space in cortical neurons (PubMed: [19906677](http://www.uniprot.org/citations/19906677)). ABPP- initiated RAGE signaling, especially stimulation of p38 mitogen- activated protein kinase (MAPK), has the capacity to drive a transport system delivering ABPP as a complex with RAGE to the intraneuronal space. Participates in endothelial albumin transcytosis together with HMGB1 through the RAGE/SRC/Caveolin-1 pathway, leading to endothelial hyperpermeability (PubMed: [27572515](http://www.uniprot.org/citations/27572515)). Mediates the loading of HMGB1 in extracellular vesicles (EVs) that shuttle HMGB1 to hepatocytes by transferrin-mediated endocytosis and subsequently promote hepatocyte pyroptosis by activating the NLRP3 inflammasome (PubMed: [34743181](http://www.uniprot.org/citations/34743181)). Promotes also extracellular hypomethylated DNA (CpG DNA) uptake by cells via the endosomal route to activate inflammatory responses (PubMed: [24081950](http://www.uniprot.org/citations/24081950), PubMed: [28515150](http://www.uniprot.org/citations/28515150)).

Cellular Location

[Isoform 1]: Cell membrane; Single-pass type I membrane protein [Isoform 10]: Cell membrane; Single-pass type I membrane protein

Tissue Location

Endothelial cells.

RAGE Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RAGE Rabbit mAb - Images