

# MMP14 Rabbit mAb

Catalog # AP78598

## Specification

## MMP14 Rabbit mAb - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW

WB, IHC-P, FC, IP, ICC <u>P50281</u> Human, Mouse, Rat Rabbit Monoclonal Antibody 65894

## MMP14 Rabbit mAb - Additional Information

Gene ID 4323

Other Names MMP14

Format Liquid

## MMP14 Rabbit mAb - Protein Information

Name MMP14

#### Function

Endopeptidase that degrades various components of the extracellular matrix such as collagen (PubMed:<a href="http://www.uniprot.org/citations/8015608" target=" blank">8015608</a>). Essential for pericellular collagenolysis and modeling of skeletal and extraskeletal connective tissues during development (By similarity). Activates progelatinase A/MMP2, thereby acting as a positive regulator of cell growth and migration (PubMed:<a href="http://www.uniprot.org/citations/22065321" target=" blank">22065321</a>, PubMed:<a href="http://www.uniprot.org/citations/8015608" target=" blank">8015608</a>). Involved in the formation of the fibrovascular tissues in association with pro-MMP2 (PubMed:<a href="http://www.uniprot.org/citations/12714657" target=" blank">12714657</a>, PubMed:<a href="http://www.uniprot.org/citations/22065321" target=" blank">22065321</a>). May be involved in actin cytoskeleton reorganization by cleaving PTK7 (PubMed:<a href="http://www.uniprot.org/citations/20837484" target="\_blank">20837484</a>). Acts as a regulator of Notch signaling by mediating cleavage and inhibition of DLL1 (PubMed:<a href="http://www.uniprot.org/citations/21572390" target=" blank">21572390</a>). Cleaves ADGRB1 to release vasculostatin-40 which inhibits angiogenesis (PubMed:<a href="http://www.uniprot.org/citations/22330140" target=" blank">22330140</a>). Acts as a negative regulator of the GDF15-GFRAL aversive response by mediating cleavage and inactivation of GFRAL (PubMed: <a href="http://www.uniprot.org/citations/35177851" target=" blank">35177851</a>).

**Cellular Location** 



Cell membrane; Single-pass type I membrane protein. Melanosome. Cytoplasm Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065). Forms a complex with BST2 and localizes to the cytoplasm (PubMed:17081065)

**Tissue Location** 

Expressed in stromal cells of colon, breast, and head and neck. Expressed in lung tumors.

#### MMP14 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

MMP14 Rabbit mAb - Images