

### CDC25C Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP7862b

### Specification

# **CDC25C Antibody (C-term) - Product Information**

Application	WB,E
Primary Accession	<u>P30307</u>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	53365
Antigen Region	376-402
, incigent region	070 101

### CDC25C Antibody (C-term) - Additional Information

Gene ID 995

Other Names M-phase inducer phosphatase 3, Dual specificity phosphatase Cdc25C, CDC25C

Target/Specificity

This CDC25C antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 376-402 amino acids from the C-terminal region of human CDC25C.

Dilution WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CDC25C Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## CDC25C Antibody (C-term) - Protein Information

Name CDC25C

**Function** Functions as a dosage-dependent inducer in mitotic control. Tyrosine protein phosphatase required for progression of the cell cycle (PubMed:<u>8119945</u>). When phosphorylated, highly effective in activating G2 cells into prophase (PubMed:<u>8119945</u>). Directly dephosphorylates



CDK1 and activates its kinase activity (PubMed:<u>8119945</u>).


Cellular Location Nucleus

## CDC25C Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

# CDC25C Antibody (C-term) - Images



Western blot analysis of anti-CDC25C Antibody (C-term) (Cat.#AP7862b) in K562 cell line lysates (35ug/lane).CDC25C(arrow) was detected using the purified Pab.

### CDC25C Antibody (C-term) - Background

CDC25C is a tyrosine phosphatase and belongs to the Cdc25 phosphatase family. It directs dephosphorylation of cyclin B-bound CDC2 and triggers entry into mitosis. It is also thought to suppress p53-induced growth arrest.

### CDC25C Antibody (C-term) - References

Varmeh,S., Mol. Cancer Ther. 7 (12), 3789-3799 (2008) Long,J.M., Cell Cycle 7 (19), 3062-3073 (2008) Bonnet,J., Cell Cycle 7 (13), 1991-1998 (2008)